

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 December 2002 (19.12.2002)

PCT

(10) International Publication Number
WO 02/100347 A2

- (51) International Patent Classification⁷: **A61K**
- (21) International Application Number: PCT/US02/18689
- (22) International Filing Date: 11 June 2002 (11.06.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/297,521 11 June 2001 (11.06.2001) US
60/298,514 14 June 2001 (14.06.2001) US
60/366,090 19 March 2002 (19.03.2002) US
- (71) Applicant: **XENOPORT, INC.** [US/US]; 3410 Central Expressway, Santa Clara, CA 95051 (US).
- (72) Inventors: **GALLOP, Mark, A.**; 511 Orange Avenue, Los Altos, CA 94022 (US). **CUNDY, Kenneth, C.**; 45 Summit Ridge Place, Redwood City, CA 94062 (US). **ZHOU, Cindy, X.**; 1054 Colorado Avenue, Palo Alto, CA 94303 (US). **YAO, Fenmei**; 2185B Stanford Avenue., Mountain View, CA 94040 (US). **XIANG, Jia-Ning**; 707 DeSoto Drive, Palo Alto, CA 94303 (US). **OLLMAN, Ian, R.**; 3222 Glendora Drive # 103, San Mateo, CA 94403 (US). **QUI, Fayang, G.**; 5 Hackett Avenue_#3, Mountain View, CA 94043 (US).
- (74) Agents: **SINGH, Sunil, K.**; Cooley Godward LLP 3000 El Camino Real, Five Palo Alto Square, Palo Alto, CA 94306-2155 et al. (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: PRODRUGS OF GABA ANALOGS, COMPOSITIONS AND USES THEREOF

(57) Abstract: The present invention provides prodrugs of GABA analogs, pharmaceutical compositions of prodrugs of GABA analogs and methods for making prodrugs of GABA 5 analogs. The present invention also provides methods for using prodrugs of GABA analogs and methods for using pharmaceutical compositions of prodrugs of GABA analogs for treating or preventing common diseases and/or disorders.

WO 02/100347 A2

PRODRUGS OF GABA ANALOGS, COMPOSITIONS AND USES THEREOF

This application claims the benefit under 35 U.S.C. § 119(e) from United States Provisional Application Serial No. 60/297,521 filed June 11, 2001; United States
5 Provisional Application Serial No. 60/298,514 filed June 14, 2001; and United States Provisional Application Serial No. 60/366,090 filed March 19, 2002, which are herein incorporated by reference.

1. Field Of The Invention


10 The present invention relates generally to prodrugs of GABA analogs, pharmaceutical compositions of to prodrugs of GABA analogs, methods of making prodrugs of GABA analogs, methods of using prodrugs of GABA analogs and pharmaceutical compositions of prodrugs of GABA analogs. More particularly, the present invention relates to prodrugs of gabapentin and pregabalin, pharmaceutical compositions of
15 prodrugs of gabapentin and pregabalin, methods of making prodrugs of gabapentin and pregabalin, methods of using prodrugs of gabapentin and pregabalin and pharmaceutical compositions of prodrugs of gabapentin and pregabalin.

2. Background Of The Invention


20 Gamma ("γ")-aminobutyric acid ("GABA") is one of the major inhibitory transmitters in the central nervous system of mammals. GABA is not transported efficiently into the brain from the bloodstream (*i.e.*, GABA does not effectively cross the blood-brain barrier). Consequently, brain cells provide virtually all of the GABA found in the brain (GABA is biosynthesized by decarboxylation of glutamic acid with pyridoxal phosphate).

25 GABA regulates neuronal excitability through binding to specific membrane proteins (*i.e.*, GABAA receptors), which results in opening of an ion channel. The entry of chloride ion through the ion channel leads to hyperpolarization of the recipient cell, which consequently prevents transmission of nerve impulses to other cells. Low levels of GABA have been observed in individuals suffering from epileptic seizures, motion disorders (*e.g.*,
30 multiple sclerosis, action tremors, tardive dyskinesia), panic, anxiety, depression, alcoholism and manic behavior.


The implication of low GABA levels in a number of common disease states and/or common medical disorders has stimulated intensive interest in preparing GABA analogs, which have superior pharmaceutical properties in comparison to GABA (*e.g.*, the ability to cross the blood brain barrier). Accordingly, a number of GABA analogs, with considerable pharmaceutical activity have been synthesized in the art (See, *e.g.*, Satzinger *et al.*, United States Patent No. 4,024,175; Silverman *et al.*, United States Patent No. 5,563,175; Horwell *et al.*, United States Patent No. 6,020,370; Silverman *et al.*, United States Patent No. 6,028,214; Horwell *et al.*, United States Patent No. 6,103,932; Silverman *et al.*, United States Patent No. 6,117,906; Silverman, International Publication No. WO 92/09560; Silverman *et al.*, International Publication No. WO 93/23383; Horwell *et al.*, International Publication No. WO 97/29101, Horwell *et al.*, International Publication No. WO 97/33858; Horwell *et al.*, International Publication No. WO 97/33859; Bryans *et al.*, International Publication No. WO 98/17627; Guglietta *et al.*, International Publication No. WO 99/08671; Bryans *et al.*, International Publication No. WO 99/21824; Bryans *et al.*, International Publication No. WO 99/31057; Belliotti *et al.*, International Publication No. WO 99/31074; Bryans *et al.*, International Publication No. WO 99/31075; Bryans *et al.*, International Publication No. WO 99/61424; Bryans *et al.*, International Publication No. WO 00/15611; Bryans, International Publication No. WO 00/31020; Bryans *et al.*,




Gabapentin
(1)



Pregabalin
(2)



Vigabatrin
(3)




Baclofen
(4)

International Publication No. WO 00/50027; and Bryans *et al.*, International Publication No. WO 02/00209).

Pharmaceutically important GABA analogs include, for example, gabapentin (1), pregabalin (2), vigabatrin (3), and baclofen (4) shown above. Gabapentin is a lipophilic GABA analog that can pass through the blood-brain barrier, which has been used to clinically treat epilepsy since 1994. Gabapentin also has potentially useful therapeutic effects in chronic pain states (*e.g.*, neuropathic pain, muscular and skeletal pain), psychiatric disorders (*e.g.*, panic, anxiety, depression, alcoholism and manic behavior), movement disorders (*e.g.*, multiple sclerosis, action tremors, tardive dyskinesia), *etc.* (Magnus, *Epilepsia*, 1999, 40:S66-S72). Currently, gabapentin is also used in the clinical management of neuropathic pain. Pregabalin, which possesses greater potency in pre-clinical models of pain and epilepsy than gabapentin is presently in Phase III clinical trials.

A significant problem with many GABA analogs is intramolecular reaction of the γ amino group with the carboxyl functionality to form the γ -lactam, as exemplified for gabapentin below. Formation of γ -lactam (5) presents serious difficulties in formulating gabapentin because of its toxicity. For example, gabapentin has a toxicity (LD_{50} , mouse) of more than 8000 mg/kg, while the corresponding lactam (5) has a toxicity (LD_{50} , mouse) of 300 mg/kg. Consequently, formation of side products such as lactams during synthesis of GABA analogs and/or formulation and/or storage of GABA analogs or compositions of GABA analogs must be minimized for safety reasons (particularly, in the case of



gabapentin).

The problem of lactam contamination of GABA analogs, particularly in the case of gabapentin, has been partially overcome through use of special additional purification steps, precise choice of adjuvant materials in pharmaceutical compositions and careful control procedures (Augurt *et al.*, United States Patent No. 6,054,482). However, attempts to prevent lactam contamination have not been entirely successful, in either synthesis or storage of GABA analogs such as gabapentin or compositions thereof.

Rapid systemic clearance is another significant problem with many GABA analogs including gabapentin, which consequently require frequent dosing to maintain a therapeutic

or prophylactic concentration in the systemic circulation (Bryans *et al.*, *Med. Res. Rev.*, 1999, 19, 149-177). For example, dosing regimens of 300-600 mg doses of gabapentin administered three times per day are typically used for anticonvulsive therapy. Higher doses (1800-3600 mg/d in divided doses) are typically used for the treatment of neuropathic
5 pain states.

Sustained released formulations are a conventional solution to the problem of rapid systemic clearance, as is well known to those of skill in the art (See, e.g., "Remington's Pharmaceutical Sciences," Philadelphia College of Pharmacy and Science, 17th Edition, 1985). Osmotic delivery systems are also recognized methods for sustained drug delivery
10 (See, e.g., Verma *et al.*, *Drug Dev. Ind. Pharm.*, 2000, 26:695-708). Many GABA analogs, including gabapentin and pregabalin, are not absorbed *via* the large intestine. Rather, these compounds are typically absorbed in the small intestine by the large neutral amino acid transporter ("LNAA") (Jezyk *et al.*, *Pharm. Res.*, 1999, 16, 519-526). The rapid passage of conventional dosage forms through the proximal absorptive region of the gastrointestinal
15 tract has prevented the successful application of sustained release technologies to many GABA analogs.

Thus, there is a significant need for effective sustained release versions of GABA analogs to minimize increased dosing frequency due to rapid systemic clearance of these compounds. There is also a need for pure GABA analogs, (particularly gabapentin and
20 pregabalin analogs) which are substantially pure and do not spontaneously lactamize during either formulation or storage.

3. Summary Of The Invention

The present invention addresses these and other needs by providing prodrugs of
25 GABA analogs, pharmaceutical compositions of prodrugs of GABA analogs and methods for making prodrugs of GABA analogs. The present invention also provides methods for using prodrugs of GABA analogs, and methods for using pharmaceutical compositions of prodrugs of GABA analogs for treating or preventing common diseases and/or disorders.

Importantly, the prodrugs provided by the present invention may possess significant
30 pharmaceutical advantages of particular use in medicine. First, the promoiety of the prodrugs of GABA analogs provided by the current invention are typically labile *in vivo* (*i.e.*, cleaved by either enzymatic or chemical means to generate substantial quantities of a GABA analog before the prodrug is cleared from a patient. Second, the promoiety derivative provided by cleavage of the promoiety from the prodrug, and any metabolite



thereof, is typically non-toxic when administered to a mammal in accordance with dosing regimens typically followed with the GABA analog.


The compounds of the instant invention have a promoiety attached to the γ amino group of GABA analogs. This promoiety may be directly attached to the γ amino group of a GABA analog, or optionally may be attached to the amino group of an α -amino acid promoiety, or to the hydroxy group of an α -hydroxy acid promoiety, which itself is attached to the γ amino group of the GABA analog.

The compounds of the invention may also have a promoiety attached to the carboxyl group of GABA analogs. The carboxyl promoiety will typically be an ester or thioester group. A wide variety of ester or thioester groups may be used to form carboxyl promoieties.

Accordingly, the compounds of the invention may include as many as four promoieties, including one carboxyl promoiety and up to three amino promoieties attached in sequence to the γ amino group (*i.e.*, such that each promoiety is sequentially cleaved from the N-terminal end of the GABA analog). The compounds of the invention may contain two amino promoieties and one carboxyl promoiety, two amino promoieties, one amino promoiety and one carboxyl promoiety or one amino promoiety. Preferably, in those compounds of the invention which contain both an amino promoiety and a carboxyl promoiety, the carboxyl promoiety is hydrolyzed prior to the complete cleavage of the promoiety (ies) attached to the amine group.

In a first aspect, the present invention provides compounds of Formula (I), Formula (II) or Formula (III):





(III)

or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:



m, n, t and u are independently 0 or 1;

X is O or NR¹⁶;

5 W is O or NR¹⁷;

Y is O or S;

R¹ is selected from the group consisting of hydrogen, R²⁴C(O)-, R²⁵OC(O)-, R²⁴C(S)-, R²⁵OC(S)-, R²⁵SC(O)-, R²⁵SC(S)-, (R⁹O)(R¹⁰O)P(O)-, R²⁵S-



10 each R² is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, substituted acyl, acylamino, substituted acylamino, alkylamino, substituted alkylamino, alkylsulfinyl, substituted alkylsulfinyl, alkylsulfonyl, substituted alkylsulfonyl, alkylthio, substituted alkylthio, alkoxycarbonyl, substituted alkoxycarbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, aryloxy, substituted aryloxy, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, dialkylamino, substituted dialkylamino, halo, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl, substituted heteroarylalkyl, heteroalkyloxy, substituted heteroalkyloxy, heteroaryloxy and substituted heteroaryloxy, or optionally, R² and R¹⁶ together with the atoms to which they are attached
15 form a cycloheteroalkyl or substituted cycloheteroalkyl ring;

20 R³ and R⁶ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl,

substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl;

R^4 and R^5 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, acyl, substituted acyl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroarylalkyl and substituted heteroarylalkyl or optionally, R^4 and R^5 together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl or bridged cycloalkyl ring;

R^8 and R^{12} are independently selected from the group consisting of hydrogen, acyl, substituted acyl, alkoxycarbonyl, substituted alkoxycarbonyl, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl, or optionally R^8 and R^{12} together with the carbon atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring;

R^{11} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, acyl, substituted acyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cyano, cycloalkyl, substituted cycloalkyl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl, substituted heteroarylalkyl, alkoxycarbonyl, substituted alkoxycarbonyl, cycloheteroalkyloxycarbonyl, substituted cycloheteroalkyloxycarbonyl, aryloxycarbonyl, substituted aryloxycarbonyl, heteroaryloxycarbonyl, substituted heteroaryloxycarbonyl and nitro;

R^7 , R^9 , R^{10} , R^{15} , R^{16} and R^{17} are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl;

R^{13} and R^{14} are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxycarbonyl, substituted alkoxycarbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloalkoxycarbonyl, substituted cycloalkoxycarbonyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl or optionally, R^{13} and R^{14} together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring;

R^{20} and R^{21} are independently selected from the group consisting of hydrogen, acyl, substituted acyl, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl or optionally R^{20} and R^{21} together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring;

R^{22} and R^{23} are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl and substituted arylalkyl or optionally, R^{22} and R^{23} together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring;

R^{24} is selected from the group consisting of hydrogen, acyl, substituted acyl, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl; and

R^{25} is selected from the group consisting of acyl, substituted acyl, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl.

In a second aspect, the present invention provides pharmaceutical compositions of compounds of the invention. The pharmaceutical compositions generally comprise one or more compounds of the invention, and a pharmaceutically acceptable vehicle.

In a third aspect, the present invention provides methods for treating or preventing epilepsy, depression, anxiety, psychosis, faintness attacks, hypokinesia, cranial disorders, neurodegenerative disorders, panic, pain (especially, neuropathic pain and muscular and skeletal pain), inflammatory disease (*i.e.*, arthritis), insomnia, gastrointestinal disorders or ethanol withdrawal syndrome. The methods generally involve administering to a patient in need of such treatment or prevention a therapeutically effective amount of a compound of the invention.


In a fourth aspect, the current invention provides pharmaceutical compositions for treating or preventing epilepsy, depression, anxiety, psychosis, faintness attacks, hypokinesia, cranial disorders, neurodegenerative disorders, panic, pain (especially,

neuropathic pain and muscular and skeletal pain), inflammatory disease (*i.e.*, arthritis), insomnia, gastrointestinal disorders or ethanol withdrawal syndrome in a patient in need of such treatment or prevention. The methods generally involve administering to a patient in need of such treatment or prevention a therapeutically effective amount of a pharmaceutical composition of the invention.

In a fifth aspect, the invention comprises a GABA analog derivative compound, M-G, for administration to a patient in need of therapy, wherein M is a promoiety and G is derived from a GABA analog, H-G (where H is hydrogen). The promoiety M, once cleaved from G, and any metabolite thereof, exhibits a carcinogenically toxic dose (TD₅₀) in rats of greater than 0.2 mmol/kg/day. Further, the promoiety M cleaves from G at a sufficient rate *in vivo*, upon colonic administration to rats, to produce:

- (i) a maximum concentration of H-G in plasma (C_{max}) of at least 120% of the C_{max} of H-G in plasma is achieved by colonically administering an equimolar dose of H-G; and
- (ii) an AUC that is at least 120 % of the AUC is achieved by colonically administering an equimolar dose of H-G.

Preferably, M-G is a derivative of Formula (XIV):




(XIV)

or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:

R is hydrogen or R and R⁶ together with the atoms to which they are attached form an azetidine, substituted azetidine, pyrrolidine or substituted pyrrolidine ring; and

Y, R³, R⁴, R⁵, R⁶ and R⁷ are as previously defined.

Most preferably, M is a derivative of Formula (XV):



(XV)

wherein:

n, X, R¹ and R² are as previously defined.

4. Detailed Description Of The Invention

5

4.1 Definitions

“Active transport or active transport process” refers to the movement of molecules across cellular membranes that:

- a) is directly or indirectly dependent on an energy mediated process (*i.e.*, driven by ATP hydrolysis, ion gradient, *etc.*);
- or
- b) occurs by facilitated diffusion mediated by interaction with specific transporter proteins.

15 “Alkyl” refers to a saturated or unsaturated, branched, straight-chain or cyclic monovalent hydrocarbon radical derived by the removal of one hydrogen atom from a single carbon atom of a parent alkane, alkene or alkyne. Typical alkyl groups include, but are not limited to, methyl; ethyls such as ethanyl, ethenyl, ethynyl; propyls such as propan-1-yl, propan-2-yl, cyclopropan-1-yl, prop-1-en-1-yl, prop-1-en-2-yl, prop-2-en-1-yl (allyl),

20 cycloprop-1-en-1-yl; cycloprop-2-en-1-yl, prop-1-yn-1-yl, prop-2-yn-1-yl, *etc.*; butyls such as butan-1-yl, butan-2-yl, 2-methyl-propan-1-yl, 2-methyl-propan-2-yl, cyclobutan-1-yl, but-1-en-1-yl, but-1-en-2-yl, 2-methyl-prop-1-en-1-yl, but-2-en-1-yl, but-2-en-2-yl, buta-1,3-dien-1-yl, buta-1,3-dien-2-yl, cyclobut-1-en-1-yl, cyclobut-1-en-3-yl, cyclobuta-1,3-dien-1-yl, but-1-yn-1-yl, but-1-yn-3-yl, but-3-yn-1-yl, *etc.*; and the like.

25 The term “alkyl” is specifically intended to include groups having any degree or level of saturation, *i.e.*, groups having exclusively single carbon-carbon bonds, groups having one or more double carbon-carbon bonds, groups having one or more triple carbon-carbon bonds and groups having mixtures of single, double and triple carbon-carbon bonds. Where a specific level of saturation is intended, the expressions “alkanyl,” “alkenyl,” and

30 “alkynyl” are used. Preferably, an alkyl group comprises from 1 to 20 carbon atoms, more preferably, from 1 to 10 carbon atoms.

“Alkanyl” refers to a saturated branched, straight-chain or cyclic alkyl radical derived by the removal of one hydrogen atom from a single carbon atom of a parent alkane.

Typical alkanyl groups include, but are not limited to, methanyl; ethanyl; propanyls such as propan-1-yl, propan-2-yl (isopropyl), cyclopropan-1-yl, etc.; butanyls such as butan-1-yl, butan-2-yl (*sec*-butyl), 2-methyl-propan-1-yl (isobutyl), 2-methyl-propan-2-yl (*t*-butyl), cyclobutan-1-yl, etc.; and the like.

5

“Alkenyl” refers to an unsaturated branched, straight-chain or cyclic alkyl radical having at least one carbon-carbon double bond derived by the removal of one hydrogen atom from a single carbon atom of a parent alkene. The group may be in either the *cis* or *trans* conformation about the double bond(s). Typical alkenyl groups include, but are not limited to, ethenyl; propenyls such as prop-1-en-1-yl, prop-1-en-2-yl, prop-2-en-1-yl (allyl), prop-2-en-2-yl, cycloprop-1-en-1-yl; cycloprop-2-en-1-yl; butenyls such as but-1-en-1-yl, but-1-en-2-yl, 2-methyl-prop-1-en-1-yl, but-2-en-1-yl, but-2-en-1-yl, but-2-en-2-yl, buta-1,3-dien-1-yl, buta-1,3-dien-2-yl, cyclobut-1-en-1-yl, cyclobut-1-en-3-yl, cyclobuta-1,3-dien-1-yl, etc.; and the like.

15

“Alkynyl” refers to an unsaturated branched, straight-chain or cyclic alkyl radical having at least one carbon-carbon triple bond derived by the removal of one hydrogen atom from a single carbon atom of a parent alkyne. Typical alkynyl groups include, but are not limited to, ethynyl; propynyls such as prop-1-yn-1-yl, prop-2-yn-1-yl, etc.; butynyls such as but-1-yn-1-yl, but-1-yn-3-yl, but-3-yn-1-yl, etc.; and the like.

20

“Acyl” refers to a radical -C(O)R, where R is hydrogen, alkyl, cycloalkyl, cycloheteroalkyl, aryl, arylalkyl, heteroalkyl, heteroaryl, heteroarylalkyl as defined herein. Representative examples include, but are not limited to formyl, acetyl, cyclohexylcarbonyl, cyclohexylmethylcarbonyl, benzoyl, benzylcarbonyl and the like.

25

“Acylamino” (or alternatively “acylamido”) refers to a radical -NR'C(O)R, where R' and R are each independently hydrogen, alkyl, cycloalkyl, cycloheteroalkyl, aryl, arylalkyl, heteroalkyl, heteroaryl, heteroarylalkyl, as defined herein. Representative examples include, but are not limited to, formylamino, acetylamino (*i.e.*, acetamido), cyclohexylcarbonylamino, cyclohexylmethyl-carbonylamino, benzoylamino (*i.e.*, benzamido), benzylcarbonylamino and the like.

30

“Acyloxy” refers to a radical -OC(O)R, where R is hydrogen, alkyl, cycloalkyl, cycloheteroalkyl, aryl, arylalkyl, heteroalkyl, heteroaryl or heteroarylalkyl, as defined herein. Representative examples include, but are not limited to, acetyloxy (or acetoxo), butyloxy (or butoxo), benzoyloxy and the like.

5

“Alkylamino” means a radical -NHR where R represents an alkyl or cycloalkyl group as defined herein. Representative examples include, but are not limited to, methylamino, ethylamino, 1-methylethylamino, cyclohexyl amino and the like.

10

“Alkoxy” refers to a radical -OR where R represents an alkyl or cycloalkyl group as defined herein. Representative examples include, but are not limited to, methoxy, ethoxy, propoxy, butoxy, cyclohexyloxy and the like.

15

“Alkoxy carbonyl” refers to a radical -C(O)-alkoxy where alkoxy is as defined herein.

20

“Alkylsulfonyl” refers to a radical -S(O)₂R where R is an alkyl or cycloalkyl group as defined herein. Representative examples include, but are not limited to, methylsulfonyl, ethylsulfonyl, propylsulfonyl, butylsulfonyl and the like.

25

“Alkylsulfinyl” refers to a radical -S(O)R where R is an alkyl or cycloalkyl group as defined herein. Representative examples include, but are not limited to, methylsulfinyl, ethylsulfinyl, propylsulfinyl, butylsulfinyl and the like.

30

“Alkylthio” refers to a radical -SR where R is an alkyl or cycloalkyl group as defined herein that may be optionally substituted as defined herein. Representative examples include, but are not limited to methylthio, ethylthio, propylthio, butylthio and the like.

“Amino” refers to the radical -NH₂.

“Aryl” refers to a monovalent aromatic hydrocarbon group derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. Typical aryl groups include, but are not limited to, groups derived from aceanthrylene,

acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, coronene, fluoranthene, fluorene, hexacene, hexaphene, hexalene, *as*-indacene, *s*-indacene, indane, indene, naphthalene, octacene, octaphene, octalene, ovalene, penta-2,4-diene, pentacene, pentalene, pentaphene, perylene, phenalene, phenanthrene, picene, pleiadene, pyrene, pyranthrene, rubicene, triphenylene, trinaphthalene and the like. Preferably, an aryl group comprises from 6 to 20 carbon atoms, more preferably between 6 to 12 carbon atoms.


“Arylalkyl” refers to an acyclic alkyl group in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with an aryl group. Typical arylalkyl groups include, but are not limited to, benzyl, 2-phenylethan-1-yl, 2-phenylethen-1-yl, naphthylmethyl, 2-naphthylethan-1-yl, 2-naphthylethen-1-yl, naphthobenzyl, 2-naphthophenylethan-1-yl and the like. Where specific alkyl moieties are intended, the nomenclature arylalkanyl, arylalkenyl and/or arylalkynyl is used. Preferably, an arylalkyl group is (C_6 - C_{30}) arylalkyl, *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the arylalkyl group is (C_1 - C_{10}) and the aryl moiety is (C_6 - C_{20}), more preferably, an arylalkyl group is (C_6 - C_{20}) arylalkyl, *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the arylalkyl group is (C_1 - C_8) and the aryl moiety is (C_6 - C_{12}).

“Arylalkyloxy” refers to an -O-arylalkyl group where arylalkyl is as defined herein.

“Aryloxycarbonyl” refers to a radical -C(O)-O-aryl where aryl is as defined herein.

“AUC” is the area under the plasma drug concentration-versus-time curve extrapolated from zero time to infinity.

“Bridged cycloalkyl” refers to a radical selected from the group consisting of



wherein:

A is $(CR^{35}R^{36})_b$;

5 R^{35} and R^{36} are independently selected from the group consisting of hydrogen and methyl;

R^{33} and R^{34} are independently selected from the group consisting of hydrogen and methyl;

b is an integer from 1 to 4; and

c is an integer from 0 to 2.

10

“Carbamoyl” refers to the radical $-C(O)N(R)_2$ where each R group is independently hydrogen, alkyl, cycloalkyl or aryl as defined herein, which may be optionally substituted, as defined herein.

15 “Carboxy” means the radical $-C(O)OH$.

“Carcinogenic potency (TD_{50})” (see Peto *et al.*, *Environmental Health Perspectives* 1984, 58, 1-8) is defined for a particular compound in a given animal species as that chronic dose-rate in mg/kg body wt/day which would induce tumors in half the test animals at the
20 end of a standard lifespan for the species. Since the tumor(s) of interest often does occur in control animals, TD_{50} is more precisely defined as: that dose-rate in mg/kg body wt/day which, if administered chronically for the standard lifespan of the species, will halve the probability of remaining tumorless throughout that period. A TD_{50} can be computed for any particular type of neoplasm, for any particular tissue, or for any combination of these.

25

“ C_{max} ” is the highest drug concentration observed in plasma following an extravascular dose of drug.

“Compounds of the invention” refers to compounds encompassed by generic
30 formulae disclosed herein and includes any specific compounds within that formula whose structure is disclosed herein. The compounds of the invention may be identified either by their chemical structure and/or chemical name. When the chemical structure and chemical name conflict, the chemical structure is determinative of the identity of the compound. The compounds of the invention may contain one or more chiral centers and/or double bonds

and therefore, may exist as stereoisomers, such as double-bond isomers (*i.e.*, geometric isomers), enantiomers or diastereomers. Accordingly, the chemical structures depicted herein encompass all possible enantiomers and stereoisomers of the illustrated compounds including the stereoisomerically pure form (*e.g.*, geometrically pure, enantiomerically pure or diastereomerically pure) and enantiomeric and stereoisomeric mixtures. Enantiomeric and stereoisomeric mixtures can be resolved into their component enantiomers or stereoisomers using separation techniques or chiral synthesis techniques well known to the skilled artisan. The compounds of the invention may also exist in several tautomeric forms including the enol form, the keto form and mixtures thereof. Accordingly, the chemical structures depicted herein encompass all possible tautomeric forms of the illustrated compounds. The compounds of the invention also include isotopically labeled compounds where one or more atoms have an atomic mass different from the atomic mass conventionally found in nature. Examples of isotopes that may be incorporated into the compounds of the invention include, but are not limited to, ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F and ^{36}Cl . Further, it should be understood, when partial structures of the compounds of the invention are illustrated, that brackets indicate the point of attachment of the partial structure to the rest of the molecule.

“Composition of the invention” refers to at least one compound of the invention and a pharmaceutically acceptable vehicle, with which the compound is administered to a patient. When administered to a patient, the compounds of the invention are administered in isolated form, which means separated from a synthetic organic reaction mixture.

“Cyano” means the radical -CN.


“Cycloalkyl” refers to a saturated or unsaturated cyclic alkyl radical. Where a specific level of saturation is intended, the nomenclature “cycloalkanyl” or “cycloalkenyl” is used. Typical cycloalkyl groups include, but are not limited to, groups derived from cyclopropane, cyclobutane, cyclopentane, cyclohexane, and the like. Preferably, the cycloalkyl group is ($\text{C}_3\text{-C}_{10}$) cycloalkyl, more preferably ($\text{C}_3\text{-C}_7$) cycloalkyl.

“Cycloheteroalkyl” refers to a saturated or unsaturated cyclic alkyl radical in which one or more carbon atoms (and any associated hydrogen atoms) are independently replaced with the same or different heteroatom. Typical heteroatoms to replace the carbon atom(s)

include, but are not limited to, N, P, O, S, Si, *etc.* Where a specific level of saturation is intended, the nomenclature “cycloheteroalkanyl” or “cycloheteroalkenyl” is used. Typical cycloheteroalkyl groups include, but are not limited to, groups derived from epoxides, imidazolidine, morpholine, piperazine, piperidine, pyrazolidine, pyrrolidine, quinuclidine, and the like.

“Cycloheteroalkyloxycarbonyl” refers to a radical -C(O)-OR where R is cycloheteroalkyl is as defined herein.

“Derived from a bile acid” refers to a moiety that is structurally related to a compound of Formulae (XVII) or (XVIII):



wherein each of D, E and F are independently H or OH.


The structure of the moiety is identical to the compounds above except at 1 or 2 positions. At these positions, a hydrogen atom attached to a hydroxyl group and/or the hydroxyl moiety of the carboxylic acid group has been replaced with a covalent bond that serves as a point of attachment to another moiety, which is preferably a GABA analog or GABA analog derivative.

“Derived from a GABA analog” refers to a moiety that is structurally related to a GABA analog. The structure of the moiety is identical to the compound except at 1 or 2 positions. At these positions, a hydrogen atom attached to the amino group, and (optionally) the hydroxyl moiety of the carboxylic acid group has been replaced with a covalent bond that serves as a point of attachment to another moiety.

“Dialkylamino” means a radical -NRR' where R and R' independently represent an alkyl or cycloalkyl group as defined herein. Representative examples include, but are not

limited to, dimethylamino, methylethylamino, di-(1-methylethyl)amino, (cyclohexyl)(methyl)amino, (cyclohexyl)(ethyl)amino, (cyclohexyl)(propyl)amino and the like.

- 5 “GABA analog” refers to a compound, unless specified otherwise, as having the following structure:



wherein:

- 10 R is hydrogen, or R and R⁶ together with the atoms to which they are attached form an azetidine, substituted azetidine, pyrrolidine or substituted pyrrolidine ring;
- R³ and R⁶ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl; and
- 15 R⁴ and R⁵ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, acyl, substituted acyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl or
- 20 optionally, R⁴ and R⁵ together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl or bridged cycloalkyl ring.

 “Halo” means fluoro, chloro, bromo, or iodo.

25

 “Heteroalkyloxy” means an -O-heteroalkyl group where heteroalkyl is as defined herein.

- “Heteroalkyl, Heteroalkanyl, Heteroalkenyl, Heteroalkynyl” refer to alkyl, alkanyl, alkenyl and alkynyl radical, respectively, in which one or more of the carbon atoms (and any associated hydrogen atoms) are each independently replaced with the same or different heteroatomic groups. Typical heteroatomic groups include, but are not limited to, -O-, -S-, -
- 30

O-O-, -S-S-, -O-S-, -NR'-, =N-N=, -N=N-, -N=N-NR'-, -PH-, -P(O)₂-, -O-P(O)₂-, -S(O)-, -S(O)₂-, -SnH₂- and the like, where R' is hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl or substituted aryl.

5 "Heteroaryl" refers to a monovalent heteroaromatic radical derived by the removal of one hydrogen atom from a single atom of a parent heteroaromatic ring system. Typical heteroaryl groups include, but are not limited to, groups derived from acridine, arsindole, carbazole, β -carboline, chromane, chromene, cinnoline, furan, imidazole, indazole, indole, indoline, indolizine, isobenzofuran, isochromene, isoindole, isoindoline, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, perimidine, phenanthridine, phenanthroline, phenazine, phthalazine, pteridine, purine, pyran, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, pyrrolizine, quinazoline, quinoline, quinolizine, quinoxaline, tetrazole, thiadiazole, thiazole, thiophene, triazole, xanthene, and the like. Preferably, the heteroaryl group is between 5-20 membered heteroaryl, more preferably
10 between 5-10 membered heteroaryl. Preferred heteroaryl groups are those derived from thiophene, pyrrole, benzothiophene, benzofuran, indole, pyridine, quinoline, imidazole, oxazole and pyrazine.

20 "Heteroaryloxycarbonyl" refers to a radical -C(O)-OR where R is heteroaryl as defined herein.

25 "Heteroarylalkyl" refers to an acyclic alkyl group in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or *sp*³ carbon atom, is replaced with a heteroaryl group. Where specific alkyl moieties are intended, the nomenclature heteroarylalkanyl, heteroarylalkenyl and/or heteroarylalkynyl is used. In preferred embodiments, the heteroarylalkyl group is a 6-30 membered heteroarylalkyl, *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the heteroarylalkyl is 1-10 membered and the heteroaryl moiety is a 5-20-membered heteroaryl, more preferably, 6-20 membered heteroarylalkyl, *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the heteroarylalkyl is 1-8
30 membered and the heteroaryl moiety is a 5-12-membered heteroaryl.

 "Passive diffusion" refers to uptake of an agent that is not mediated by a specific transporter protein. An agent that is substantially incapable of passive diffusion has a

permeability across a standard cell monolayer (*e.g.*, Caco-2) *in vitro* of less than 5×10^{-6} cm/sec, and usually less than 1×10^{-6} cm/sec (in the absence of an efflux mechanism).

5 “Pharmaceutically acceptable” means approved or approvable by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopoeia or other generally recognized pharmacopoeia for use in animals, and more particularly in humans.

10 “Pharmaceutically acceptable salt” refers to a salt of a compound of the invention, which is pharmaceutically acceptable and possesses the desired pharmacological activity of the parent compound. Such salts include: (1) acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-
15 (4-hydroxybenzoyl) benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic acid, glucoheptonic acid, 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic
20 acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like; or (2) salts formed when an acidic proton present in the parent compound is replaced by a metal ion, *e.g.*, an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, N-methylglucamine and the like.

25

 “Pharmaceutically acceptable vehicle” refers to a diluent, adjuvant, excipient or carrier with which a compound of the invention is administered.

30 “Patient” includes humans. The terms “human” and “patient” are used interchangeably herein.

 “Preventing” or “prevention” refers to a reduction in risk of acquiring a disease or disorder (*i.e.*, causing at least one of the clinical symptoms of the disease not to develop in a

patient that may be exposed to or predisposed to the disease but does not yet experience or display symptoms of the disease).

“Prodrug” refers to a derivative of a drug molecule that requires a transformation
 5 within the body to release the active drug. Prodrugs are frequently (though not necessarily) pharmacologically inactive until converted to the parent drug.

“Promoiety” refers to a form of protecting group that when used to mask a functional group within a drug molecule converts the drug into a prodrug. Typically, the
 10 promoiety will be attached to the drug *via* bond(s) that are cleaved by enzymatic or non-enzymatic means *in vivo*.

“Protecting group” refers to a grouping of atoms that when attached to a reactive functional group in a molecule masks, reduces or prevents reactivity of the functional group.
 15 Examples of protecting groups can be found in Green *et al.*, “Protective Groups in Organic Chemistry”, (Wiley, 2nd ed. 1991) and Harrison *et al.*, “Compendium of Synthetic Organic Methods”, Vols. 1-8 (John Wiley and Sons, 1971-1996). Representative amino protecting groups include, but are not limited to, formyl, acetyl, trifluoroacetyl, benzyl, benzyloxycarbonyl (“CBZ”), *tert*-butoxycarbonyl (“Boc”), trimethylsilyl (“TMS”), 2-
 20 trimethylsilyl-ethanesulfonyl (“SES”), trityl and substituted trityl groups, allyloxycarbonyl, 9-fluorenylmethyloxycarbonyl (“FMOC”), nitro-veratryloxycarbonyl (“NVOC”) and the like. Representative hydroxy protecting groups include, but are not limited to, those where the hydroxy group is either acylated or alkylated such as benzyl, and trityl ethers as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers and allyl ethers.

25

“Substituted” refers to a group in which one or more hydrogen atoms are each independently replaced with the same or different substituent(s). Typical substituents include, but are not limited to, -X, -R²⁹, -O⁻, =O, -OR²⁹, -SR²⁹, -S⁻, =S, -NR²⁹R³⁰, =NR²⁹, -CX₃, -CF₃, -CN, -OCN, -SCN, -NO, -NO₂, =N₂, -N₃, -S(O)₂O⁻, -S(O)₂OH, -S(O)₂R²⁹, -
 30 OS(O₂)O⁻, -OS(O)₂R²⁹, -P(O)(O⁻)₂, -P(O)(OR²⁹)(O⁻), -OP(O)(OR²⁹)(OR³⁰), -C(O)R²⁹, -C(S)R²⁹, -C(O)OR²⁹, -C(O)NR²⁹R³⁰, -C(O)O⁻, -C(S)OR²⁹, -NR³¹C(O)NR²⁹R³⁰, -NR³¹C(S)NR²⁹R³⁰, -NR³¹C(NR²⁹)NR²⁹R³⁰ and -C(NR²⁹)NR²⁹R³⁰, where each X is independently a halogen; each R²⁹ and R³⁰ are independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted

cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl, substituted heteroarylalkyl, -NR³¹R³², -C(O)R³¹ or -S(O)₂R³¹ or optionally R²⁹ and R³⁰ together with the atom to which they are both attached form a cycloheteroalkyl or substituted cycloheteroalkyl ring; and R³¹ and R³² are independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl or substituted heteroarylalkyl.

10 “Transporter protein” refers to a protein that has a direct or indirect role in transporting a molecule into and/or through a cell. For example, a transporter protein may be, but is not limited to, solute carrier transporters, co- transporters, counter transporters, uniporters, symporters, antiporters, pumps, equilibrative transporters, concentrative transporters and other proteins, which mediate active transport, energy-dependent transport, facilitated diffusion, exchange mechanisms and specific absorption mechanisms.

15 Transporter proteins, may also be, but are not limited to, membrane-bound proteins that recognize a substrate and effect its entry into or exit from a cell by a carrier-mediated transporter or by receptor-mediated transport. A transporter protein, may also be, but is not limited to, an intracellularly expressed protein that participates in trafficking of substrates through or out of a cell. Transporter proteins, may also be, but are not limited to, proteins or glycoproteins exposed on the surface of a cell that do not directly transport a substrate but bind to the substrate holding it in proximity to a receptor or transporter protein that effects entry of the substrate into or through the cell. Examples of carrier proteins include: the intestinal and liver bile acid transporters, dipeptide transporters, oligopeptide

20 transporters, simple sugar transporters (e.g., SGLT1), phosphate transporters, monocarboxylic acid transporters, P-glycoprotein transporters, organic anion transporters (OAT), and organic cation transporters. Examples of receptor-mediated transport proteins include: viral receptors, immunoglobulin receptors, bacterial toxin receptors, plant lectin receptors, bacterial adhesion receptors, vitamin transporters and cytokine growth factor

25 receptors.

30 receptors.

 “Treating” or “treatment” of any disease or disorder refers, in one embodiment, to ameliorating the disease or disorder (*i.e.*, arresting or reducing the development of the disease or at least one of the clinical symptoms thereof). In another embodiment “treating”

or "treatment" refers to ameliorating at least one physical parameter, which may not be discernible by the patient. In yet another embodiment, "treating" or "treatment" refers to inhibiting the disease or disorder, either physically, (*e.g.*, stabilization of a discernible symptom), physiologically, (*e.g.*, stabilization of a physical parameter), or both. In yet another embodiment, "treating" or "treatment" refers to delaying the onset of the disease or disorder.


"Therapeutically effective amount" means the amount of a compound that, when administered to a patient for treating a disease, is sufficient to effect such treatment for the disease. The "therapeutically effective amount" will vary depending on the compound, the disease and its severity and the age, weight, *etc.*, of the patient to be treated.


Reference will now be made in detail to preferred embodiments of the invention. While the invention will be described in conjunction with the preferred embodiments, it will be understood that it is not intended to limit the invention to those preferred embodiments. To the contrary, it is intended to cover alternatives, modifications, and equivalents as may be included within the spirit and scope of the invention as defined by the appended claims.

4.2 The Compounds of the Invention


Those of skill in the art will appreciate that compounds of Formulae (I), (II) and (III) share certain structural features in common. These compounds are all GABA analogs (*i.e.*, γ -aminobutyric acid derivatives) to which promoieties have been attached. In particular, R^2 , R^3 , R^4 , R^5 , R^6 , X and Y are common substituents found in compounds of Formulae (I), (II) and (III).

The compounds of the invention include compounds of Formula (I), Formula (II) or Formula (III):





(II)



(III)

or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:

n , t , u , X , Y , R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^{20} , R^{21} , R^{22} and R^{23} are as previously

5 defined above.

In a preferred embodiment, compounds of Formulae (I), (II) and (III) do not include the following compounds:

when R^3 and R^6 are both hydrogen, then R^4 and R^5 are not both hydrogen and are not both methyl;

10 in a compound of Formula (I) when either n is 0 or when n is 1 and X is NR^{16} , then R^1 is not hydrogen;

in a compound of Formula (I) neither R^1 , R^7O- , $R^{24}C(O)-$, $R^{25}C(O)-$ nor $R^{25}O-$ is a moiety derived from a bile acid;

15 in a compound of Formula (I) when R^1 is $R^{24}C(O)-$ and n is 0, then R^{24} is not methyl, *tert*-butyl, 2-aminoethyl, 3-aminopropyl, benzyl, phenyl or 2-(benzoyloxymethyl)phenyl;

20 in a compound of Formula (I) when R^1 is $R^{25}OC(O)-$, then R^{25} is not $R^{26}C(O)CR^{13}R^{14}-$, wherein R^{26} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl;

in a compound of Formula (I) when R^1 is $R^{25}OC(O)-$ and n is 0, then R^{25} is not methyl, *tert*-butyl or benzyl;

in a compound of Formula (I) when n is 0 and R^1 is $R^{25}C(O)OCR^{13}R^{14}OC(O)-$ then if either R^{13} or R^{14} is hydrogen, alkoxy carbonyl, substituted alkoxy carbonyl, carbamoyl, cycloalkoxy carbonyl or substituted cycloalkoxy carbonyl, the other of R^{13} or R^{14} is not hydrogen;

- 5 in a compound of Formula (I) when n is 1, X is NH , R^3 , R^5 and R^6 are each hydrogen, and R^4 is cyclohexyl, then R^2 is not benzyl;

in a compound of Formula (II) when t is 1, u is 0, then neither R^{20} nor R^{21} is 2-hydroxy-3-methyl-5-chlorophenyl; and

in a compound of Formula (II) when u is 1 and X is O , then t is 1.


- 10 In one embodiment of compounds of Formulae (I), (II) and (III), when R^3 and R^6 are each hydrogen, then R^4 and R^5 are neither both hydrogen nor both methyl.

- In one embodiment of compounds of Formula (I), when either n is 0 or when n is 1 and X is NR^{16} , then R^1 is not hydrogen. In another embodiment, of compounds of Formula (I), neither R^1 , R^7O- , $R^{24}C(O)-$, $R^{25}C(O)-$ nor $R^{25}O-$ is a moiety derived from a bile acid. In still
- 15 another embodiment of compounds of Formula (I), when R^1 is $R^{24}C(O)-$ and n is 0, then R^{24} is not alkyl, substituted alkyl, arylalkyl, aryl or substituted aryl. In still another embodiment of compounds of Formula (I), when R^1 is $R^{24}C(O)-$ and n is 0, then R^{24} is not C_{1-4} alkanyl, benzyl, phenyl or substituted phenyl. In still another embodiment of compounds of Formula (I), when R^1 is $R^{24}C(O)-$ and n is 0, then R^{24} is not methyl, *tert*-butyl, 2-
- 20 aminoethyl, 3-aminopropyl, benzyl, phenyl or 2-(benzoyloxymethyl)-phenyl. In still another embodiment of compounds of Formula (I) when R^1 is $R^{25}OC(O)-$, then R^{25} is not $R^{26}C(O)CR^{13}R^{14}-$. In still another embodiment of compounds of Formula (I) when R^1 is $R^{25}OC(O)-$ and n is 0, then R^{25} is not alkyl or arylalkyl. In still another embodiment of compounds of Formula (I) when R^1 is $R^{25}OC(O)-$ and n is 0, then R^{25} is not C_{1-4} alkanyl or
- 25 benzyl. In still another embodiment of compounds of Formula (I) when R^1 is $R^{25}OC(O)-$ and n is 0, then R^{25} is not methyl, *tert*-butyl or benzyl. In still another embodiment of compounds of Formula (I) when n is 0 and R^1 is $R^{25}C(O)OCR^{13}R^{14}OC(O)-$ then if either R^{13} or R^{14} is hydrogen, alkoxy carbonyl, substituted alkoxy carbonyl, carbamoyl, cycloalkoxy carbonyl or substituted cycloalkoxy carbonyl, the other of R^{13} or R^{14} is not
- 30 hydrogen. In still another embodiment of compounds of Formula (I) when R^3 , R^5 and R^6 are each hydrogen, then R^4 is not cyclohexyl. In still another embodiment of compounds of Formula (I) when n is 1, X is NH , R^3 , R^5 , R^6 are each hydrogen and R^2 is benzyl, then R^4 is not cyclohexyl.

In one embodiment of compounds of Formula (II), neither R^{20} nor R^{21} is 2-hydroxy-3-methyl-5-chlorophenyl. In one embodiment of compounds of Formula (II), when u is 1 and X is O, then t is 1.

In one embodiment of compounds of Formulae, (I) (II) and (III), n is 0. In another embodiment, n is 1. When n is 1, and X is NR^{16} , preferably the α -amino acid is of the L-stereochemical configuration.

In another embodiment of compounds of Formulae (I) and (II), R^7 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkanyl, substituted arylalkanyl, cycloalkanyl, substituted cycloalkanyl, cycloheteroalkanyl and substituted cycloheteroalkanyl. In a preferred embodiment, Y is O and R^7 is hydrogen. In still another embodiment, Y is O and R^7 is alkanyl, substituted alkanyl, alkenyl, substituted alkenyl, aryl or substituted aryl. Preferably, R^7 is methyl, ethyl, benzyl, $-C(CH_3)=CH_2$, $-CH_2C(O)N(CH_3)_2$,



where V is O or CH_2 .

In one preferred embodiment of compounds of Formulae (I), (II) and (III), R^2 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl. Preferably, R^2 is selected from the group consisting hydrogen, alkanyl, substituted alkanyl, aryl, substituted aryl, arylalkanyl, substituted arylalkanyl, cycloalkanyl, heteroarylalkyl and substituted heteroarylalkanyl.

In another embodiment compounds of Formulae (I), (II) and (III), X is NH and R^2 is hydrogen, cycloalkanyl or alkanyl. Preferably, R^2 is hydrogen, methyl, isopropyl, isobutyl, *sec*-butyl, *t*-butyl, cyclopentyl or cyclohexyl. In another embodiment, X is NH and R^2 is substituted alkanyl. Preferably, R^2 is $-CH_2OH$, $-CH(OH)CH_3$, $-CH_2CO_2H$, $-CH_2CH_2CO_2H$, $-CH_2CONH_2$, $-CH_2CH_2CONH_2$, $-CH_2CH_2SCH_3$, CH_2SH , $-CH_2(CH_2)_3NH_2$ or $-CH_2CH_2CH_2NHC(NH)NH_2$. In still another embodiment, X is NH and R^2 is selected from the group consisting of aryl, arylalkanyl, substituted arylalkanyl and heteroarylalkanyl. Preferably, R^2 is phenyl, benzyl, 4-hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl or 2-indolyl. In yet another embodiment, X is NR^{16} and R^2 and R^{16} together with the atoms to which they are attached form a cycloheteroalkyl or substituted cycloheteroalkyl ring.

Preferably R^2 and R^{16} together with the atoms to which they are attached form an azetidine, pyrrolidine or piperidine ring.

In still another embodiment of compounds of Formulae (I), (II) and (III), R^3 is hydrogen. In still another embodiment, R^6 is hydrogen. In yet another embodiment, R^3 and R^6 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl and substituted cycloalkyl. Preferably, R^3 and R^6 are independently selected from the group consisting of hydrogen and alkanyl. More preferably, R^3 is hydrogen or alkanyl and R^6 is hydrogen.


In still another preferred embodiment of compounds of Formulae (I), (II) and (III), R^4 and R^5 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl and substituted cycloheteroalkyl. Preferably, R^4 and R^5 are independently selected from the group consisting of hydrogen, alkanyl and substituted alkanyl.

In another embodiment of compounds of Formulae (I), (II) and (III), R^4 and R^5 together with the carbon atom to which they are attached form a cycloalkanyl or substituted cycloalkanyl ring. Preferably, R^4 and R^5 together with the carbon atom to which they are attached form a cyclobutyl, substituted cyclobutyl, cyclopentyl, substituted cyclopentyl, cyclohexyl or substituted cyclohexyl ring. In another embodiment, R^4 and R^5 together with the carbon atom to which they are attached form a cycloheteroalkyl or substituted cycloheteroalkyl ring. In still another embodiment, R^4 and R^5 together with the carbon atom to which they are attached form a bridged cycloalkyl ring.

In one embodiment of compounds of Formula (I), n is 1, R^1 is $R^{24}C(O)-$ or $R^{24}C(S)-$ and R^{24} is alkyl, substituted alkyl, heteroalkyl, substituted heteroalkyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl. Preferably, R^{24} is methyl, ethyl, 2-propyl, t-butyl, - $CH_2OCH(CH_3)_2$, phenyl or 3-pyridyl.

In another embodiment of compounds of Formula (I), n is 1, R^1 is $R^{25}OC(O)-$ or $R^{25}SC(O)-$ and R^{25} is alkyl, substituted alkyl, heteroalkyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl. Preferably, R^{25} is ethyl, 2-propyl, neopentyl, $-CH_2OCH(CH_3)_2$, phenyl or 2-pyridyl.

One preferred embodiment of compounds of Formula (I) includes compounds of Formula (IV):



or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:

n , Y , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^{13} , R^{14} , R^{16} and R^{25} are as previously defined.

In a preferred embodiment, the compounds of Formula (IV), do not include the
 5 following compounds:

when either R^{13} or R^{14} is hydrogen, alkoxycarbonyl, substituted alkoxycarbonyl, carbamoyl, cycloalkoxycarbonyl or substituted cycloalkoxycarbonyl, then the other of R^{13} or R^{14} is not hydrogen; and

$R^{25}C(O)$ is not a moiety derived from a bile acid.

10 In one embodiment of compounds of Formulae (IV), R^{13} and R^{14} are independently hydrogen, alkyl, substituted alkyl, alkoxycarbonyl, aryl, arylalkyl, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloalkoxycarbonyl or heteroaryl (preferably, when R^{13} is alkoxycarbonyl, cycloalkoxycarbonyl or carbamoyl then R^{14} is methyl). More preferably, R^{13} and R^{14} are independently hydrogen, methyl, ethyl, propyl, isopropyl, butyl, isobutyl,
 15 *sec*-butyl, *tert*-butyl, cyclopentyl, cyclohexyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, *sec*-butoxycarbonyl, *tert*-butoxycarbonyl, cyclohexyloxycarbonyl, phenyl, benzyl, phenethyl or 3-pyridyl.

In another embodiment of compounds of Formula (IV), R^{13} and R^{14} are
 20 independently hydrogen, alkanyl, substituted alkanyl, cycloalkanyl or substituted cycloalkanyl. Preferably, R^{13} and R^{14} are hydrogen, alkanyl or cycloalkanyl. More preferably, R^{13} and R^{14} are independently hydrogen, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, *tert*-butyl, cyclopentyl or cyclohexyl. Even more preferably, R^{13} is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, *tert*-butyl, cyclopentyl or
 25 cyclohexyl and R^{14} is hydrogen, or R^{13} is methyl and R^{14} is methyl.

In still another embodiment of compounds of Formula (IV), R^{13} and R^{14} are independently hydrogen, aryl, arylalkyl or heteroaryl. More preferably, R^{13} and R^{14} are independently hydrogen, phenyl, benzyl, phenethyl or 3-pyridyl. Even more preferably, R^{13} is phenyl, benzyl, phenethyl or 3-pyridyl and R^{14} is hydrogen.

30 In still another embodiment of compounds of Formula (IV), R^{13} and R^{14} are independently hydrogen, alkyl, substituted alkyl, alkoxycarbonyl, carbamoyl, or

cycloalkoxycarbonyl. Preferably, when R¹³ is alkoxycarbonyl, cycloalkoxycarbonyl or carbamoyl then R¹⁴ is methyl. More preferably, R¹³ is methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, *sec*-butoxycarbonyl, *tert*-butoxycarbonyl or cyclohexyloxycarbonyl and R¹⁴ is methyl.

- 5 In still another embodiment of compounds of Formula (IV), R¹³ and R¹⁴ together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring. Preferably, R¹³ and R¹⁴ together with the carbon atom to which they are attached form a cycloalkyl ring. More preferably, R¹³ and R¹⁴ together with the carbon atom to which they are attached form a cyclobutyl, 10 cyclopentyl or cyclohexyl ring.

- In still another embodiment of compounds of Formula (IV), R²⁵ is acyl, substituted acyl, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl or substituted 15 heteroarylalkyl. Preferably, R²⁵ is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl. More preferably, R²⁵ is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1- 20 dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, 25 cyclohexyl or 3-pyridyl.

In still another embodiment of compounds of Formula (IV), R²⁵ is acyl or substituted acyl. More preferably, R²⁵ is acetyl, propionyl, butyryl, benzoyl or phenacetyl.

- In still another embodiment of compounds of Formula (IV), R²⁵ is alkanyl or substituted alkanyl. Preferably, R²⁵ is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1- 30 (1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-

yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl or 1-(1,3-dioxan-2-yl)-2-phenethyl. More preferably, R²⁵ is methyl, ethyl, propyl, isopropyl, butyl, 1,1-dimethoxyethyl or 1,1-diethoxyethyl.

In still another embodiment of compounds of Formula (IV), R²⁵ is aryl, arylalkyl or heteroaryl. Preferably, R²⁵ is phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl or 3-pyridyl.

In still another embodiment of compounds of Formula (IV), R²⁵ is cycloalkyl or substituted cycloalkyl. More preferably R²⁵ is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

In still another embodiment of compounds of Formula (IV), R²⁵ is acyl, substituted acyl, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl or substituted heteroarylalkyl; and R¹³ and R¹⁴ are independently hydrogen, alkyl, substituted alkyl, alkoxycarbonyl, substituted alkoxycarbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloalkoxycarbonyl, substituted cycloalkoxycarbonyl, heteroaryl or substituted heteroaryl (preferably, when R¹³ is alkoxycarbonyl, substituted alkoxycarbonyl, cycloalkoxycarbonyl, substituted cycloalkoxycarbonyl or carbamoyl then R¹⁴ is methyl). Preferably, R²⁵ is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl and R¹³ and R¹⁴ are independently hydrogen, alkyl, substituted alkyl, alkoxycarbonyl, aryl, arylalkyl, carbamoyl, cycloalkyl, cycloalkoxycarbonyl or heteroaryl (preferably, when R¹³ is alkoxycarbonyl, substituted alkoxycarbonyl, cycloalkoxycarbonyl, substituted cycloalkoxycarbonyl or carbamoyl then R¹⁴ is methyl). More preferably, R²⁵ is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or 3-pyridyl and R¹³ and R¹⁴ are independently hydrogen, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, *tert*-

butyl, cyclopentyl, cyclohexyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, *sec*-butoxycarbonyl, *tert*-butoxycarbonyl, cyclohexyloxycarbonyl, phenyl, benzyl, phenethyl or 3-pyridyl. Even more preferably, R²⁵ is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, cyclohexyl or 3-pyridyl, and R¹³ and R¹⁴ are independently hydrogen, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, *tert*-butyl, cyclopentyl, cyclohexyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, cyclohexyloxycarbonyl, phenyl, benzyl, phenethyl or 3-pyridyl.

In still another embodiment of compounds of Formula (IV), R²⁵ is acyl, substituted acyl, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl or substituted heteroarylalkyl and R¹³ and R¹⁴ together with the atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring. Preferably, R²⁵ is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl and R¹³ and R¹⁴ together with the atom to which they are attached form a cycloalkyl or substituted cycloalkyl ring. More preferably, R²⁵ is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or 3-pyridyl, and R¹³ and R¹⁴ together with the atom to which they are attached form a cyclobutyl, cyclopentyl or a cyclohexyl ring.

In still another embodiment of compounds of Formula (IV), R²⁵ is acyl or substituted acyl and R¹³ and R¹⁴ are independently hydrogen, alkyl, substituted alkyl, alkoxycarbonyl, substituted alkoxycarbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloalkoxycarbonyl, substituted cycloalkoxycarbonyl, heteroaryl or substituted heteroaryl (preferably, when R¹³ is

alkoxycarbonyl, substituted alkoxycarbonyl, cycloalkoxycarbonyl, substituted cycloalkoxycarbonyl or carbamoyl then R¹⁴ is methyl). Preferably, R²⁵ is acetyl, propionyl, butyryl, benzoyl or phenacetyl, and R¹³ and R¹⁴ are independently hydrogen, alkyl, substituted alkyl, alkoxycarbonyl, substituted alkoxycarbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloalkoxycarbonyl, substituted cycloalkoxycarbonyl, heteroaryl or substituted heteroaryl (preferably, when R¹³ is alkoxycarbonyl, cycloalkoxycarbonyl or carbamoyl then R¹⁴ is methyl).

In still another embodiment of compounds of Formula (IV), R²⁵ is alkanyl or substituted alkanyl and R¹³ and R¹⁴ are independently hydrogen, alkyl, substituted alkyl, alkoxycarbonyl, substituted alkoxycarbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloalkoxycarbonyl, substituted cycloalkoxycarbonyl, heteroaryl or substituted heteroaryl (preferably, when R¹³ is alkoxycarbonyl, substituted alkoxycarbonyl, cycloalkoxycarbonyl, substituted cycloalkoxycarbonyl or carbamoyl then R¹⁴ is methyl). Preferably, R²⁵ is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl or 1-(1,3-dioxan-2-yl)-2-phenethyl, and R¹³ and R¹⁴ are independently hydrogen, alkyl, substituted alkyl, alkoxycarbonyl, substituted alkoxycarbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloalkoxycarbonyl, substituted cycloalkoxycarbonyl, heteroaryl or substituted heteroaryl (preferably, when R¹³ is alkoxycarbonyl, cycloalkoxycarbonyl or carbamoyl then R¹⁴ is methyl).

In still another embodiment of compounds of Formula (IV), R²⁵ is aryl, substituted aryl, arylalkyl, substituted arylalkyl, heteroaryl or substituted heteroaryl, and R¹³ and R¹⁴ are independently hydrogen, alkyl, substituted alkyl, alkoxycarbonyl, substituted alkoxycarbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloalkoxycarbonyl, substituted cycloalkoxycarbonyl, heteroaryl or substituted heteroaryl (preferably, when R¹³ is alkoxycarbonyl, substituted alkoxycarbonyl, cycloalkoxycarbonyl, substituted cycloalkoxycarbonyl or carbamoyl then

R¹⁴ is methyl). Preferably R²⁵ is phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl or 3-pyridyl and R¹³ and R¹⁴ are independently hydrogen, alkyl, substituted alkyl, alkoxy carbonyl, substituted alkoxy carbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloalkoxy carbonyl, substituted cycloalkoxy carbonyl, heteroaryl or substituted heteroaryl (preferably, when R¹³ is alkoxy carbonyl, cycloalkoxy carbonyl or carbamoyl then R¹⁴ is methyl).

In still another embodiment of compounds of Formula (IV), R²⁵ is cycloalkyl or substituted cycloalkyl, and R¹³ and R¹⁴ are independently hydrogen, alkyl, substituted alkyl, alkoxy carbonyl, substituted alkoxy carbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloalkoxy carbonyl, substituted cycloalkoxy carbonyl, heteroaryl or substituted heteroaryl (preferably, when R¹³ is alkoxy carbonyl, substituted alkoxy carbonyl, cycloalkoxy carbonyl, substituted cycloalkoxy carbonyl or carbamoyl then R¹⁴ is methyl). Preferably, R²⁵ is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl and R¹³ and R¹⁴ are independently hydrogen, alkyl, substituted alkyl, alkoxy carbonyl, substituted alkoxy carbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloalkoxy carbonyl, substituted cycloalkoxy carbonyl, heteroaryl or substituted heteroaryl (preferably, when R¹³ is alkoxy carbonyl, cycloalkoxy carbonyl or carbamoyl then R¹⁴ is methyl).

In still another embodiment of compounds of Formula (IV), R²⁵ is acyl, substituted acyl, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl or substituted heteroarylalkyl, and R¹³ and R¹⁴ are independently hydrogen, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl. Preferably, R²⁵ is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl and R¹³ and R¹⁴ are independently hydrogen, alkyl, substituted alkyl, cycloalkyl or substituted cycloalkyl. More preferably, R²⁵ is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl and R¹³ and R¹⁴ are independently hydrogen, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, *tert*-butyl, cyclopentyl or cyclohexyl. In the above embodiments, R²⁵ is preferably methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-

dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or 3-pyridyl.

In still another embodiment of compounds of Formula (IV), R^{25} is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl and R^{13} and R^{14} are independently hydrogen, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl.

10 Preferably, R^{25} is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl and R^{13} and R^{14} are independently hydrogen, aryl, arylalkyl or heteroaryl. More preferably, R^{25} is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl and R^{13} and R^{14} are independently hydrogen, phenyl, benzyl, phenethyl or 3-pyridyl. In the above embodiments, R^{25} is preferably methyl, ethyl, propyl, isopropyl,

15 butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl,

20 phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or 3-pyridyl.


In still another embodiment of compounds of Formula (IV), R^{25} is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl and R^{13} and R^{14} are independently hydrogen, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl.

25 Preferably, R^{25} is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl and R^{13} and R^{14} are independently hydrogen, alkyl, substituted alkyl, alkoxycarbonyl, substituted alkoxycarbonyl, carbamoyl, cycloalkoxycarbonyl or substituted cycloalkoxycarbonyl (preferably, when R^{13} is alkoxycarbonyl, substituted alkoxycarbonyl, carbamoyl, cycloalkoxycarbonyl or substituted cycloalkoxycarbonyl then R^{14} is methyl; more preferably, R^{13} is methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, *sec*-butoxycarbonyl, *tert*-butoxycarbonyl or cyclohexyloxycarbonyl, and R^{14} is methyl). In the above embodiments,

30

- R^{25} is preferably methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or 3-pyridyl.
- 10 In still another embodiment of compounds of Formula (IV), R^{25} is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl and R^{13} and R^{14} together with the atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring. Preferably, R^{25} is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl and R^{13} and R^{14}
- 15 together with the atom to which they are attached form a cycloalkyl or substituted cycloalkyl ring. More preferably R^{25} is acyl, substituted acyl, alkyl, substituted alkyl, aryl, arylalkyl, cycloalkyl or heteroaryl, and R^{13} and R^{14} together with the atom to which they are attached form a cyclobutyl, cyclopentyl or cyclohexyl ring. In the above embodiments, R^{25} is preferably methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or 3-pyridyl.

In still another embodiment of compounds of Formulae (I) and (III), R^1 is



- 30 m is 0, and R^8 , R^{11} and R^{12} are as previously defined.

In one embodiment of compounds of Formulae (I) and (III), R^{11} is acyl, alkoxy carbonyl, aryloxy carbonyl, cycloalkoxy carbonyl or carbamoyl, R^8 is hydrogen,

alkoxycarbonyl, alkyl, aryl, arylalkyl or cyano and R^{12} is hydrogen, alkoxycarbonyl, alkyl, substituted alkyl, aryl, or arylalkyl.

In another embodiment of compounds of Formulae (I) and (III), R^{11} is selected from the group consisting of acetyl, propionyl, butyryl, isobutyryl, pivaloyl, cyclopentanecarbonyl, cyclohexanecarbonyl, benzoyl, phenacetyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, *sec*-butoxycarbonyl, *tert*-butoxycarbonyl, cyclopentylloxycarbonyl, cyclohexylloxycarbonyl, phenoxycarbonyl, benzyloxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N-propylcarbamoyl, N-isopropylcarbamoyl, N-butylcarbamoyl, N-isobutylcarbamoyl, N-*sec*-butylcarbamoyl, N-*tert*-butylcarbamoyl, N-cyclopentylcarbamoyl, N-cyclohexylcarbamoyl, N-phenylcarbamoyl, N-benzylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, N,N-dipropylcarbamoyl, N,N-diisopropylcarbamoyl, N,N-dibutylcarbamoyl, N,N-dibenzylcarbamoyl, N-pyrrolidinylcarbamoyl, N-piperidinylcarbamoyl and N-morpholinylcarbamoyl. More preferably, R^{11} is selected from the group consisting of acetyl, propionyl, butyryl, isobutyryl, cyclohexanecarbonyl, benzoyl, phenacetyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, cyclohexylloxycarbonyl, phenoxycarbonyl, benzyloxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N-propylcarbamoyl, N-isopropylcarbamoyl, N-phenylcarbamoyl, N-benzylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, N,N-dipropylcarbamoyl, N-pyrrolidinylcarbamoyl, N-piperidinylcarbamoyl and N-morpholinylcarbamoyl.

In still another embodiment of compounds of Formulae (I) and (III), R^8 is selected from the group consisting of hydrogen, methyl, ethyl, propyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, phenoxycarbonyl, benzyloxycarbonyl and cyano. More preferably R^8 is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl and butoxycarbonyl.

In still another embodiment of compounds of Formulae (I) and (III), R^{12} is selected from the group consisting of hydrogen, methyl, ethyl, propyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, phenoxycarbonyl and benzyloxycarbonyl. More preferably R^{12} is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl and butoxycarbonyl.

In still another embodiment of compounds of Formulae (I) and (III), R^{11} is selected from the group consisting of hydrogen, alkoxycarbonyl, alkyl, substituted alkyl, aryl, arylalkyl and R^8 and R^{12} together with the carbon atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring.

5 Preferably, R^{11} is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl and butoxycarbonyl, and R^8 and R^{12} together with the carbon atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring. More preferably, R^{11} is hydrogen or methyl and R^8 and R^{12} together with the carbon atoms to which they are


10 attached form a cyclopent-1-ene, cyclohex-1-ene, 2-cyclopenten-1-one, 2-cyclohexen-1-one, 2-(5H)-furanone or 5,6-dihydro-pyran-2-one ring.

In still another embodiment of compounds of Formulae (I) and (III), R^{12} is selected from the group consisting of hydrogen, alkoxycarbonyl, alkyl, substituted alkyl, aryl, arylalkyl and R^8 and R^{11} together with the carbon atoms to which they are attached form a

15 cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring. Preferably, R^{12} is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl and butoxycarbonyl, and R^8 and R^{11} together with the carbon atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring. More preferably, R^{12} is


20 selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl and butoxycarbonyl, and R^8 and R^{11} together with the carbon atoms to which they are attached form a γ -butyrolactone, δ -valerolactone or 2,2-dimethyl-1,3-dioxan-4,6-dione ring.

In still another embodiment of compounds of Formulae (I) and (III), R^1 is



25 and R^{15} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl. Preferably, R^{15} is methyl, ethyl, propyl, isopropyl, cyclopentyl, cyclohexyl, phenyl, 4-hydroxyphenyl, benzyl, 4-hydroxybenzyl or 3-pyridyl.

30 In still another embodiment, of Formulae (I) and (III), R^1 is




where R^{37} is hydrogen, alkyl, substituted alkyl, acyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, heterocycloalkyl, substituted cycloheteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl or substituted heteroarylalkyl;

5 Z is O, N or S; and

Ar is aryl, substituted aryl, heteroaryl or substituted heteroaryl.

Preferably, Z and $\text{CH}_2\text{OC(O)-}$ are in a conjugated relationship with one another (e.g., 1,4 or 1,2 related in a six membered ring system).

In still another embodiment, of Formulas (I) and (III), R^1 is



10

where q is 0 or 1;

R^{38} and R^{39} are independently hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted alkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl;

15 R^{40} and R^{41} are independently hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted alkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl; or together with the carbon atom to which they are attached form a cycloalkyl ring

R^{42} and R^{43} are independently alkyl, substituted alkyl, cycloalkyl, substituted alkyl, aryl, substituted aryl, heteroaryl and substituted aryl or together with the carbon atoms to which they are attached form an aryl, substituted aryl, heteroaryl or substituted aryl ring;

20 and


R^{37} is as previously defined.

In a preferred embodiment of compounds of Formulae (I) - (IV), Y is O, R^3 , R^6 and R^7 are hydrogen and R^4 and R^5 together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, bridged cycloalkyl or substituted bridged cycloalkyl ring. In another preferred embodiment

of compounds of Formulae (I) - (IV), R^4 and R^5 together with the carbon atom to which they are attached form a cycloalkyl or substituted cycloalkyl ring. In one embodiment n is 0, t is 0 and u is 0. In another embodiment, n is 1 and R^2 is hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *t*-butyl, cyclopentyl, cyclohexyl, phenyl, benzyl, 4-hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl, 2-indolyl, $-\text{CH}_2\text{OH}$, $-\text{CH}(\text{OH})\text{CH}_3$, $-\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{SCH}_3$, CH_2SH , $-\text{CH}_2(\text{CH}_2)_3\text{NH}_2$ or $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NHC}(\text{NH})\text{NH}_2$. In still another embodiment, n is 1 and R^2 and R^{16} together with the atoms to which they are attached form a pyrrolidine ring.


In still another preferred embodiment of compounds of Formulae (I) - (IV), R^4 and R^5 together with the carbon atom to which they are attached form a cyclobutyl or substituted cyclobutyl ring. Preferably, the substituted cyclobutyl ring is substituted with one or more substituents selected from the group consisting of alkanyl, substituted alkanyl, halo, hydroxy, carboxyl and alkoxycarbonyl.

In still another preferred embodiment of compounds of Formulae (I) - (IV), R^4 and R^5 together with the carbon atom to which they are attached form a cyclopentyl or substituted cyclopentyl ring. Preferably, the cyclopentyl ring is substituted with alkanyl, substituted alkanyl, halo, hydroxy, carboxyl or alkoxycarbonyl. More preferably, the cyclopentyl ring is substituted with alkanyl. Even more preferably, the cyclopentyl ring is selected from the group consisting of



Preferably, in a more specific version of the above embodiments, R^7 is hydrogen.


In still another preferred embodiment of compounds of Formulae (I) - (IV), R^4 and R^5 together with the carbon atom to which they are attached form a cyclohexyl or substituted cyclohexyl ring. Preferably, the cyclohexyl ring is substituted with alkanyl, substituted alkanyl, halo, hydroxy, carboxyl or alkoxycarbonyl. More preferably, the cyclohexyl ring is substituted with alkanyl. Even more preferably, the cyclohexyl ring is selected from the group consisting of



Preferably, in a more specific version of the above embodiments, R^7 is hydrogen.

- 5 In still another preferred embodiment of compounds of Formulae (I) - (IV), R^4 and R^5 together with the carbon atom to which they are attached form a cycloheteroalkyl or substituted cycloheteroalkyl ring.

In one embodiment, n is 0. In another embodiment, n is 1, and R^2 is hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *t*-butyl, cyclopentyl, cyclohexyl, phenyl, benzyl, 4-




- 10 hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl, 2-indolyl, $-\text{CH}_2\text{OH}$, $-\text{CH}(\text{OH})\text{CH}_3$, $-\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{SCH}_3$, CH_2SH , $-\text{CH}_2(\text{CH}_2)_3\text{NH}_2$ or $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NHC}(\text{NH})\text{NH}_2$. In still another embodiment, n is 1 and R^2 and R^{16} together with the atoms to which they are attached form a pyrrolidine ring.
- 15 Preferably, R^4 and R^5 together with the carbon atom to which they are attached form a cycloheteroalkanyl ring. More preferably, the cycloheteroalkanyl ring is selected from the group consisting of

wherein Z is O, $\text{S}(\text{O})_p$ or NR^{18} ;

p is 0, 1 or 2; and


R^{18} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, acyl and alkoxy carbonyl. More preferably, the cycloheteroalkanyl ring is selected from the group consisting of



Preferably, in a more specific version of the above embodiments, R^7 is hydrogen.

- 5 In still another embodiment of compounds of Formulae (I) - (IV), R^4 and R^5 together with the carbon atom to which they are attached form a bridged cycloalkyl ring. In one embodiment, n is 0. In another embodiment, n is 1 and R^2 is hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *t*-butyl, cyclopentyl, cyclohexyl, phenyl, benzyl, 4-hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl, 2-indolyl, $-\text{CH}_2\text{OH}$, $-\text{CH}(\text{OH})\text{CH}_3$, $-\text{CH}_2\text{CO}_2\text{H}$, -
- 10 $\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{SCH}_3$, CH_2SH , $-\text{CH}_2(\text{CH}_2)_3\text{NH}_2$ or $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NHC}(\text{NH})\text{NH}_2$. In another embodiment, n is 1 and R^2 and R^{16} together with the atoms to which they are attached form a pyrrolidine ring. Preferably, the bridged cycloalkyl group is

15




Preferably, in a more specific version of the above embodiments, R^7 is hydrogen.

- In still another embodiment of compounds of Formulae (I) - (IV), Y is O, R^6 and R^7 are hydrogen, R^4 is alkyl or cycloalkyl, R^5 is hydrogen or alkyl and R^3 is hydrogen or alkyl. In one embodiment, n is 0. In another embodiment, n is 1 and R^2 is hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *t*-butyl, cyclopentyl, cyclohexyl, phenyl, benzyl, 4-hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl, 2-indolyl, $-\text{CH}_2\text{OH}$, $-\text{CH}(\text{OH})\text{CH}_3$, $-\text{CH}_2\text{CO}_2\text{H}$, -
- 20 $\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{SCH}_3$, CH_2SH , $-\text{CH}_2(\text{CH}_2)_3\text{NH}_2$

or $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NHC}(\text{NH})\text{NH}_2$. In still another embodiment, n is 1 and R^2 and R^{16} together with the atoms to which they are attached form a pyrrolidine ring. Preferably, R^4 is cycloalkyl, R^5 is hydrogen or methyl, and R^3 is hydrogen or methyl. Preferably, R^3 is hydrogen, R^4 is isobutyl and R^5 is hydrogen.

- 5 In still another embodiment of compounds of Formulae (I) - (IV), Y is O, R^5 and R^7 are hydrogen or alkanyl, R^3 and R^6 are hydrogen and R^4 is substituted heteroalkyl. Preferably, R^4 is



A is NR^{19} , O or S;

- B is alkyl, substituted alkyl, alkoxy, halogen, hydroxy, carboxyl, alkoxycarbonyl or
10 amino;

R^{19} is hydrogen, alkyl, cycloalkyl or aryl;

j is an integer from 0 to 4;


k is an integer from 1 to 4; and

l is an integer from 0 to 3.

- 15 More preferably, k is 1.


In still another embodiment of compounds of Formulae (I) - (IV), Y is O, R^5 and R^7 are hydrogen or alkanyl, R^3 and R^6 are hydrogen and R^4 is substituted alkanyl, cycloalkanyl or substituted cycloalkanyl. Preferably, R^4 is selected from the group consisting of

Preferably, R^4 is




- 20 h is an integer from 1 to 6; and
 i is an integer from 0 to 6.

More preferably, h is 1, 2, 3 or 4 and i is 0 or 1. Even more preferably, R⁴ is selected from the group consisting of



Preferably, compounds of Formulae (I) - (IV) are derived from a GABA analog of Formula (XIII):





(XIII)

wherein the GABA analog of Formula (XIII) is selected from the group consisting of:

- 1-Aminomethyl-1-cyclohexane acetic acid;
- 1-Aminomethyl-1-(3-methylcyclohexane) acetic acid;
- 1-Aminomethyl-1-(4-methylcyclohexane) acetic acid;
- 1-Aminomethyl-1-(4-isopropylcyclohexane) acetic acid;
- 1-Aminomethyl-1-(4-*tert*-butylcyclohexane) acetic acid;
- 1-Aminomethyl-1-(3,3-dimethylcyclohexane) acetic acid;
- 1-Aminomethyl-1-(3,3,5,5-tetramethylcyclohexane) acetic acid;
- 1-Aminomethyl-1-cyclopentane acetic acid;
- 1-Aminomethyl-1-(3-methylcyclopentane) acetic acid;
- 1-Aminomethyl-1-(3,4-dimethylcyclopentane) acetic acid;
- 7-Aminomethyl-bicyclo[2.2.1]hept-7-yl acetic acid;
- 9-Aminomethyl-bicyclo[3.3.1]non-9-yl acetic acid;
- 4-Aminomethyl-4-(tetrahydropyran-4-yl) acetic acid;
- 3-Aminomethyl-3-(tetrahydropyran-3-yl) acetic acid;
- 4-Aminomethyl-4-(tetrahydrothiopyran-4-yl) acetic acid;
- 3-Aminomethyl-3-(tetrahydrothiopyran-3-yl) acetic acid;
- 3-Aminomethyl-5-methyl-hexanoic acid;
- 3-Aminomethyl-5-methyl-heptanoic acid;
- 3-Aminomethyl-5-methyl-octanoic acid;
- 3-Aminomethyl-5-methyl-nonanoic acid;


- 3-Aminomethyl-5-methyl-decanoic acid;
 3-Aminomethyl-5-cyclopropyl-hexanoic acid;
 3-Aminomethyl-5-cyclobutyl-hexanoic acid;
 3-Aminomethyl-5-cyclopentyl-hexanoic acid;
 5 3-Aminomethyl-5-cyclohexyl-hexanoic acid;
 3-Aminomethyl-5-phenyl-hexanoic acid;
 3-Aminomethyl-5-phenyl-pentanoic acid;
 3-Aminomethyl-4-cyclobutyl-butyric acid;
 3-Aminomethyl-4-cyclopentyl-butyric acid;
 10 3-Aminomethyl-4-cyclohexyl-butyric acid;
 3-Aminomethyl-4-phenoxy-butyric acid;
 3-Aminomethyl-5-phenoxy-hexanoic acid; and
 3-Aminomethyl-5-benzylsulfanyl-pentanoic acid.
- 15 Particularly preferred embodiments of Formula (I) include compounds of Formulae (V) and (VI):



where R^1 , R^2 , R^7 and R^{16} are as previously defined.

- In one embodiment of compounds of Formulae (V) and (VI), n is 0. In another embodiment, n is 1 and R^2 is hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *t*-butyl, cyclopentyl, cyclohexyl, phenyl, benzyl, 4-hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl, 2-indolyl, $-\text{CH}_2\text{OH}$, $-\text{CH}(\text{OH})\text{CH}_3$, $-\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{SCH}_3$, $-\text{CH}_2\text{SH}$, $-\text{CH}_2(\text{CH}_2)_3\text{NH}_2$ or $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NHC}(\text{NH})\text{NH}_2$. Preferably, in the above embodiments, R^7 is hydrogen.
- 20


In another embodiment of compounds of Formulae (V) and (VI), n is 1, R¹ is R²⁴C(O)- or R²⁴C(S)-; and R²⁴ is alkyl, substituted alkyl, heteroalkyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl. Preferably, R²⁴ is methyl, ethyl, 2-propyl, t-butyl, -CH₂OCH(CH₃)₂, phenyl or 3-pyridyl. Preferably, in this embodiment, R⁷ is hydrogen, alkanyl, substituted alkanyl, alkenyl, substituted alkenyl, aryl or substituted aryl. More preferably, R⁷ is hydrogen, methyl, ethyl, benzyl, -C(CH₃)=CH₂, -CH₂C(O)N(CH₃)₂,



where V is O or CH₂.

Most preferably, R⁷ is hydrogen.


10 In still another embodiment of compounds of Formulae (V) and (VI), n is 1, R¹ is R²⁵OC(O)- or R²⁵SC(O)-; and R²⁵ is alkyl, substituted alkyl, heteroalkyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl. Preferably, R²⁵ is ethyl, 2-propyl, neopentyl, -CH₂OCH(CH₃)₂, phenyl or 2-pyridyl. Preferably in this embodiment, R⁷ is hydrogen, alkanyl, substituted alkanyl, alkenyl, substituted alkenyl, aryl or substituted aryl. More preferably, R⁷ is hydrogen, methyl, ethyl, benzyl, -C(CH₃)=CH₂, -CH₂C(O)N(CH₃)₂,



where V is O or CH₂.

Most preferably, R⁷ is hydrogen.


In still another embodiment of compounds of Formulae (V) and (VI), R¹ is



20

and R¹⁵ is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl. Preferably, R¹⁵ is methyl, ethyl, propyl, isopropyl, cyclopentyl, cyclohexyl, phenyl, 4-hydroxyphenyl, benzyl, 4-hydroxybenzyl or 3-pyridyl. In a more specific version of this embodiment, R⁷ is hydrogen, alkanyl, substituted alkanyl, alkenyl, substituted alkenyl, aryl or substituted aryl. More preferably, R⁷ is hydrogen, methyl, ethyl, benzyl, -C(CH₃)=CH₂, -CH₂C(O)N(CH₃)₂,

25



where V is O or CH₂.


Preferably, R⁷ is hydrogen.

Particularly preferred embodiments of compounds of Formulae (V) and (VI) are
 5 compounds selected from the group consisting of:

1-[[[(5-methyl-2-oxo-1,3-dioxol-4-en-4-yl)methoxy]carbonyl]aminomethyl]-1-cyclohexane acetic acid; and

3-[[[(5-methyl-2-oxo-1,3-dioxol-4-en-4-yl)methoxy]carbonyl]aminomethyl]-5-methyl-hexanoic acid.

10 In still another embodiment of compounds of Formulae (V) and (VI), R¹ is



m is 0, and R⁸, R¹¹ and R¹² are as previously defined. In one embodiment of compounds of Formulae (V) and (VI), R¹¹ is acyl, alkoxycarbonyl, aryloxy carbonyl, cycloalkoxycarbonyl, carbamoyl or substituted carbamoyl, R⁸ is hydrogen, alkoxycarbonyl, alkyl, aryl, arylalkyl
 15 or cyano and R¹² is hydrogen, alkoxycarbonyl, alkyl, substituted alkyl, aryl, or arylalkyl. In another embodiment of compounds of Formulae (V) and (VI), R¹¹ is selected from the group consisting of acetyl, propionyl, butyryl, isobutyryl, pivaloyl, cyclopentanecarbonyl, cyclohexanecarbonyl, benzoyl, phenacetyl, methoxycarbonyl, ethoxycarbonyl,
 20 propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, *sec*-butoxycarbonyl, *tert*-butoxycarbonyl, cyclopentylloxycarbonyl, cyclohexylloxycarbonyl, phenoxycarbonyl, benzyloxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N-propylcarbamoyl, N-isopropylcarbamoyl, N-butylcarbamoyl, N-isobutylcarbamoyl, N-*sec*-butylcarbamoyl, N-*tert*-butylcarbamoyl, N-cyclopentylcarbamoyl, N-cyclohexylcarbamoyl, N-phenylcarbamoyl, N-benzylcarbamoyl, N,N-dimethylcarbamoyl,
 25 N,N-diethylcarbamoyl, N,N-dipropylcarbamoyl, N,N-diisopropylcarbamoyl, N,N-dibutylcarbamoyl, N,N-dibenzylcarbamoyl, N-pyrrolidinylcarbamoyl, N-piperidinylcarbamoyl and N-morpholinylcarbamoyl. In still another embodiment of compounds of Formulae (V) and (VI), R¹¹ is selected from the group consisting of acetyl,
 30 propionyl, butyryl, isobutyryl, cyclohexanecarbonyl, benzoyl, phenacetyl,

methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, cyclohexyloxycarbonyl, phenoxycarbonyl, benzyloxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N-propylcarbamoyl, N-isopropylcarbamoyl, N-phenylcarbamoyl, N-benzylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, 5 N,N-dipropylcarbamoyl, N-pyrrolidinylcarbamoyl, N-piperidinylcarbamoyl and N-morpholinylcarbamoyl.

In one embodiment of compounds of Formulae (V) and (VI), R⁸ is selected from the group consisting of hydrogen, methyl, ethyl, propyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, 10 phenoxycarbonyl, benzyloxycarbonyl and cyano. Preferably, R⁸ is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl and butoxycarbonyl.


In still another embodiment of compounds of Formulae (V) and (VI), R¹² is selected from the group consisting of hydrogen, methyl, ethyl, propyl, isopropyl, phenyl, benzyl, 15 methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, phenoxycarbonyl and benzyloxycarbonyl. Preferably R¹² is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl and butoxycarbonyl.

In still another embodiment of compounds of Formulae (V) and (VI), R¹¹ is selected 20 from the group consisting of hydrogen, alkoxycarbonyl, alkyl, substituted alkyl, aryl, arylalkyl and R⁸ and R¹² together with the carbon atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring. Preferably, R¹¹ is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl and butoxycarbonyl, and R⁸ and R¹² 25 together with the carbon atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring. More preferably, R¹¹ is hydrogen or methyl and R⁸ and R¹² together with the carbon atoms to which they are attached form a 2-cyclopenten-1-one, 2-cyclohexen-1-one, 2-(5H)-furanone or 5,6-dihydro-pyran-2-one ring.

30 In still another embodiment of compounds of Formulae (V) and (VI), R¹² is selected from the group consisting of hydrogen, alkoxycarbonyl, alkyl, substituted alkyl, aryl, arylalkyl and R⁸ and R¹¹ together with the carbon atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring. Preferably, R¹² is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl,

phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl and butoxycarbonyl, and R^8 and R^{11} together with the carbon atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring. More preferably, R^{12} is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl and butoxycarbonyl, and R^8 and R^{11} together with the carbon atoms to which they are attached form a γ -butyrolactone, δ -valerolactone or 2,2-dimethyl-1,3-dioxan-4,6-dione ring.

In a more specific version of the above embodiments of compounds of Formulae (V) and (VI), R^7 is hydrogen, alkanyl, substituted alkanyl, alkenyl, substituted alkenyl, aryl or substituted aryl. More preferably, R^7 is hydrogen, methyl, ethyl, benzyl, $-C(CH_3)=CH_2$, $-CH_2C(O)N(CH_3)_2$,



where V is O or CH_2 .

Most preferably R^7 is hydrogen.

Particularly preferred embodiments of compounds of Formulae (V) and (VI) are compounds selected from the group consisting of:



Piperidinium 1-((1-methyl-3-oxo-but-1-enyl)aminomethyl)-1-cyclohexane acetate;

Piperidinium 1-{1-[(2-oxo-tetrahydrofuran-3-ylidene)ethyl]aminomethyl}-1-cyclohexane acetate;

Piperidinium 1-((2-carbomethoxy-cyclopent-1-enyl)aminomethyl)-1-cyclohexane acetate; and

Piperidinium 1-((1-methyl-2-(ethoxycarbonyl)-3-ethoxy-3-oxoprop-1-enyl)aminomethyl)-1-cyclohexane acetate.

In a particularly preferred embodiment, compounds of Formula (IV) have the structure of Formulae (VII) or (VIII):



or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:

$n, R^2, R^7, R^{13}, R^{14}, R^{16}$ and R^{25} are as previously defined.

In one preferred embodiment, the compounds of Formulae (VII) and (VIII) do not
5 include the following compounds:


if either R¹³ or R¹⁴ is hydrogen, alkoxy carbonyl, substituted alkoxy carbonyl, carbamoyl, cycloalkoxy carbonyl or substituted cycloalkoxy carbonyl, then the other of R¹³ or R¹⁴ is not hydrogen; and

R²⁵C(O) is not a moiety derived from a bile acid.

10 In one embodiment of compounds of Formulae (VII) and (VIII), n is 0. In another embodiment, n is 1. When n is 1, preferably the α -amino acid is of the L-stereochemical configuration.

In another embodiment of compounds of Formulae (VII) and (VIII), R⁷ is hydrogen, alkanyl, substituted alkanyl, alkenyl, substituted alkenyl, aryl or substituted aryl.

15 Preferably, R⁷ is hydrogen, methyl, ethyl, benzyl, -C(CH₃)=CH₂, -CH₂C(O)N(CH₃)₂,



where V is O or CH₂.

Most preferably, R⁷ is hydrogen.

20 In still another embodiment of compounds of Formulae (VII) and (VIII), n is 1, R¹⁶ is hydrogen and R² is hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *tert*-butyl, cyclopentyl, cyclohexyl, phenyl, benzyl, 4-hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl, 2-indolyl, -CH₂OH, -CH(OH)CH₃, -CH₂CO₂H, -CH₂CH₂CO₂H, -CH₂CONH₂, -CH₂CH₂CONH₂, -

$\text{CH}_2\text{CH}_2\text{SCH}_3$, $-\text{CH}_2\text{SH}$, $-\text{CH}_2(\text{CH}_2)_3\text{NH}_2$ or $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NHC}(\text{NH})\text{NH}_2$. Preferably R^{16} is hydrogen and R^2 is hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *tert*-butyl, cyclohexyl, phenyl or benzyl. In still another embodiment, n is 1 and R^2 and R^{16} together with the atoms to which they are attached form a pyrrolidine ring.

5 In still another embodiment of compounds of Formulae (VII) and (VIII), R^{25} is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R^{13} is methyl and R^{14} is hydrogen.

10 In still another embodiment of compounds of Formulae (VII) and (VIII), R^{25} is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R^{13} is ethyl and R^{14} is hydrogen.

25 In still another embodiment of compounds of Formulae (VII) and (VIII), R^{25} is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R^{13} is ethyl and R^{14} is hydrogen.

phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is propyl and R¹⁴ is hydrogen.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is
 5 selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl,
 10 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is isopropyl and R¹⁴ is hydrogen.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is
 15 selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl,
 20 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is butyl and R¹⁴ is hydrogen.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is
 selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl,
 30 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl,

phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is isobutyl and R¹⁴ is hydrogen.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is *sec*-butyl and R¹⁴ is hydrogen.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is *tert*-butyl and R¹⁴ is hydrogen.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl,

phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is cyclopentyl and R¹⁴ is hydrogen.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is cyclohexyl and R¹⁴ is hydrogen.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is methyl and R¹⁴ is methyl.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl,

phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is methoxycarbonyl and R¹⁴ is methyl.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is ethoxycarbonyl and R¹⁴ is methyl.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is propoxycarbonyl and R¹⁴ is methyl.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl,

phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is isopropoxycarbonyl and R¹⁴ is methyl.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is butoxycarbonyl and R¹⁴ is methyl.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is isobutoxycarbonyl and R¹⁴ is methyl.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is isobutoxycarbonyl and R¹⁴ is methyl.

phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is *sec*-butoxycarbonyl and R¹⁴ is methyl.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is *tert*-butoxycarbonyl and R¹⁴ is methyl.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is cyclohexyloxycarbonyl and R¹⁴ is methyl.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl,

phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is phenyl and R¹⁴ is hydrogen.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is benzyl and R¹⁴ is hydrogen.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is phenethyl and R¹⁴ is hydrogen.

In still another embodiment of compounds of Formulae (VII) and (VIII), R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl,

phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl, R¹³ is 3-pyridyl and R¹⁴ is hydrogen.

Particularly preferred embodiments of compounds of Formulae (VII) and (VIII) include compounds selected from the group consisting of:

- 5 1-{{{(α -Acetoxyethoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Propanoyloxyethoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Butanoyloxyethoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Isobutanoyloxyethoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Pivaloxyethoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 10 1-{{{(α -Benzoyloxyethoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Acetoxybutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Butanoyloxybutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Isobutanoyloxybutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Benzoyloxybutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 15 1-{{{(α -Acetoxyisobutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Propanoyloxyisobutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Butanoyloxyisobutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Isobutanoyloxyisobutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Pivaloxyisobutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 20 1-{{{(α -2,2-Diethoxypropanoyloxyisobutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -2-(1,3-Dioxolan-2-yl)propanoyloxyisobutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -(2-Amino-2-methylpropanoyl)oxyisobutoxy)carbonyl]aminomethyl}-1-
- 25 cyclohexane acetic acid;
- 1-{{{(α -Benzoyloxyisobutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Nicotinoyloxyisobutoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Acetoxyisopropoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Butanoyloxyisopropoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 30 1-{{{(α -Isobutanoyloxyisopropoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Benzoyloxyisopropoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;
- 1-{{{(α -Acetoxybenzyloxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;

1-{{[(α -Benzoyloxybenzyloxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;

1-{{[(1-(3-Methylbutanoyloxy)-2-phenylethoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;

1-{{[(1-Benzoyloxy-2-phenylethoxy)carbonyl]aminomethyl}-1-cyclohexane acetic acid;

1-{{[N-[(α -Isobutanoyloxyethoxy)carbonyl]-4-bromophenylalaninyl]-aminomethyl}-1-cyclohexane acetic acid;

3-{{[(α -Isobutanoyloxyethoxy)carbonyl]aminomethyl}-5-methylhexanoic acid;


3-{{[(α -Isobutanoyloxyisobutoxy)carbonyl]aminomethyl}-5-methylhexanoic acid;

10 and

3-{{[(α -Benzoyloxyisobutoxy)carbonyl]aminomethyl}-5-methylhexanoic acid.

In one embodiment, the compounds of the invention have the structure of Formula

(II):




(II)


15

In an embodiment of compounds of Formula (II), when R^3 , R^5 and R^6 are hydrogen, R^4 is not phenyl or substituted phenyl. More preferably, R^4 is not 4-chlorophenyl.

In a preferred embodiment, the compounds of Formula (II) have the structure of Formulae (IX) and (X):



(IX)




(X)

20

In one embodiment of compounds of Formulae (IX) and (X), t is 0. In another embodiment, t is 1 and R^2 is hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *t*-butyl, cyclopentyl, cyclohexyl, phenyl, benzyl, 4-hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl, 2-indolyl, $-\text{CH}_2\text{OH}$, $-\text{CH}(\text{OH})\text{CH}_3$, $-\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CONH}_2$, -

- 5 $\text{CH}_2\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{SCH}_3$, $-\text{CH}_2\text{SH}$, $-\text{CH}_2(\text{CH}_2)_3\text{NH}_2$ or - $\text{CH}_2\text{CH}_2\text{CH}_2\text{NHC}(\text{NH})\text{NH}_2$.


- In still another embodiment of compounds of Formulae (IX) or (X), R^{20} and R^{21} are independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl. Preferably, R^{20} and R^{21} are independently
 10 selected from the group consisting of alkyl, substituted aryl and heteroaryl. In one embodiment, R^{20} is methyl and R^{21} is methyl. Preferably, in this last embodiment, R^7 is hydrogen, methyl, ethyl, benzyl, $-\text{C}(\text{CH}_3)=\text{CH}_2$, $-\text{CH}_2\text{C}(\text{O})\text{N}(\text{CH}_3)_2$,



where V is O or CH_2 .

- 15 Most preferably, R^7 is hydrogen.


- In another embodiment of compounds of Formulae (IX) or (X), R^{20} and R^{21} together with the carbon atom to which they are attached form a cycloalkyl or substituted cycloalkyl ring. In one embodiment, R^{20} and R^{21} together with the carbon atom to which they are attached form a cyclohexyl ring. Preferably, in this last embodiment, R^7 is hydrogen,
 20 methyl, ethyl, benzyl, $-\text{C}(\text{CH}_3)=\text{CH}_2$, $-\text{CH}_2\text{C}(\text{O})\text{N}(\text{CH}_3)_2$,



where V is O or CH_2 .


Most preferably, R^7 is hydrogen.

- In one embodiment, the compounds of the invention have the structure of Formula
 25 (III):



In one embodiment of the compounds of Formula (III), n is 1, R^1 is hydrogen and R^2 is arylalkyl. Preferably, R^2 is benzyl. In another embodiment of the compounds of Formula (III), n is 0 and R^1 is $R^{25}OC(O)-$. Preferably, R^{25} is alkyl or substituted alkyl. More preferably, R^{25} is ethyl. In still another embodiment of the compounds of Formula (III), R^{22} and R^{23} are hydrogen. In still another embodiment, R^{22} and R^{23} are alkyl or substituted alkyl. Preferably, R^{22} and R^{23} are methyl.


In a preferred embodiment, the compounds of Formula (III) have the structure of Formula (XI):



(XI)

In one embodiment of the compounds of Formula (XI), n is 1, X is NH , Y is O , R^1 is hydrogen, R^2 is benzyl, R^{22} is methyl and R^{23} are methyl. In another embodiment, n is 0, Y is O , R^1 is $R^{25}OC(O)-$, R^{25} is ethyl, R^{22} is hydrogen and R^{23} is hydrogen.

In another embodiment, the compounds of Formula (III) have the structure of Formula (XII):



(XII)


In one embodiment of the compounds of Formula (XII), n is 1, X is NH , Y is O , R^1 is hydrogen, R^2 is benzyl, R^{22} is methyl and R^{23} are methyl. In another embodiment, n is 0, Y is O , R^1 is $R^{25}OC(O)-$, R^{25} is ethyl, R^{22} is hydrogen and R^{23} is hydrogen.

The instant invention also comprises a GABA analog derivative, M-G, for administration to a patient in need of therapy, wherein M is a promoiety and G is derived from a GABA analog, H-G (where H is hydrogen). The promoiety M , once cleaved from G , and any metabolite thereof, exhibits a carcinogenically toxic dose (TD_{50}) in rats of

greater than 0.2 mmol/kg/day. Further, the promoiety M is cleaved from G at a sufficient rate *in vivo*, upon colonic administration to rats, to produce:

- (i) a maximum concentration of H-G in plasma (C_{\max}) of at least 120% of the C_{\max} of H-G in plasma achieved by colonically administering an equimolar dose of H-G; and
- (ii) an AUC that is at least 120 % of the AUC achieved by colonically administering an equimolar dose of H-G.

Preferably, M-G has the structure of Formula (XIV):



(XIV)

or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:

M is a promoiety;

Y is O or S;


R is hydrogen, or R and R^6 together with the atoms to which they are attached form an azetidine, substituted azetidine, pyrrolidine or substituted pyrrolidine ring;

R^3 and R^6 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl;

R^4 and R^5 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, acyl, substituted acyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl or optionally, R^4 and R^5 together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl or bridged cycloalkyl ring; and

R^7 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl.


In a preferred embodiment, M has the structure of Formula (XV):



wherein:

n, X, R¹ and R² are as previously defined

- 5 In one embodiment, M-G include compounds wherein H-G, once cleaved from M, is substantially free of any lactam having the structure of Formula (XVI):



wherein R is hydrogen and R³, R⁴, R⁵ and R⁶ are as previously defined

- Preferably, promoiety M or any metabolite formed from M does not form
- 10 formaldehyde or pivalic acid upon cleavage of G-M. In one embodiment, the promoiety M is cleaved from G at a sufficient rate *in vivo*, upon colonic administration to rats, to produce a C_{max} of H-G in plasma of at least 200%, and most preferably at least 1000%, of the C_{max} of H-G in plasma achieved by colonically administering an equimolar dose of H-G.
- Preferably, the promoiety M is cleaved from G at a sufficient rate *in vivo*, upon colonic
- 15 administration to rats, to produce an AUC of H-G in plasma of at least 200%, and most preferably at least 500%, of the AUC of H-G in plasma achieved by colonically administering an equimolar dose of H-G. In another embodiment, the promoiety M is cleaved from H-G at a sufficient rate *in vivo*, following oral administration to dogs (*e.g.*, using an osmotic mini-pump device) at a dose of about 60 μmol equivalent of H-G per kg,
- 20 to produce a plasma concentration of H-G at 12 hours post dosing of at least 200% of the plasma concentration of H-G achieved from an equimolar dose of H-G following the same mode of administration.

4.3 Synthesis of The Compounds of the Invention

- 25 The compounds of the invention may be obtained *via* the synthetic methods illustrated in Schemes 1-17. Those of skill in the art will appreciate that a preferred synthetic route to the compounds of the invention consists of attaching promoieties to


GABA analogs. Numerous methods have been described in the art for the synthesis of GABA analogs (See, *e.g.*, Satzinger *et al.*, United States Patent No. 4,024,175; Silverman *et al.*, United States Patent No. 5,563,175; Horwell *et al.*, United States Patent No. 6,020,370; Silverman *et al.*, United States Patent No. 6,028,214; Horwell *et al.*, United States Patent
5 No. 6,103,932; Silverman *et al.*, United States Patent No. 6,117,906; Silverman, International Publication No. WO 92/09560; Silverman *et al.*, International Publication No. WO 93/23383; Horwell *et al.*, International Publication No. WO 97/29101, Horwell *et al.*, International Publication No. WO 97/33858; Horwell *et al.*, International Publication No. WO 97/33859; Bryans *et al.*, International Publication No. WO 98/17627; Guglietta *et al.*,
10 International Publication No. WO 99/08671; Bryans *et al.*, International Publication No. WO 99/21824; Bryans *et al.*, International Publication No. WO 99/31057; Belliotti *et al.*, International Publication No. WO 99/31074; Bryans *et al.*, International Publication No. WO 99/31075; Bryans *et al.*, International Publication No. WO 99/61424; Bryans *et al.*, International Publication No. WO 00/15611; Bryans, International Publication No. WO
15 00/31020; and Bryans *et al.*, International Publication No. WO 00/50027). Other methods are known in the art for synthesizing GABA analogs, which are readily accessible to the skilled artisan. The promoieties described herein, are known in the art and may be prepared and attached to GABA analogs by established procedures (See *e.g.*, Green *et al.*, "Protective Groups in Organic Chemistry", (Wiley, 2nd ed. 1991); Harrison *et al.*, "Compendium of
20 Synthetic Organic Methods", Vols. 1-8 (John Wiley and Sons, 1971-1996); "Beilstein Handbook of Organic Chemistry," Beilstein Institute of Organic Chemistry, Frankfurt, Germany; Feiser *et al.*, "Reagents for Organic Synthesis," Volumes 1-17, Wiley Interscience; Trost *et al.*, "Comprehensive Organic Synthesis," Pergamon Press, 1991; "Theilheimer's Synthetic Methods of Organic Chemistry," Volumes 1-45, Karger, 1991;
25 March, "Advanced Organic Chemistry," Wiley Interscience, 1991; Larock "Comprehensive Organic Transformations," VCH Publishers, 1989; Paquette, "Encyclopedia of Reagents for Organic Synthesis," John Wiley & Sons, 1995, Bodanzsky, "Principles of Peptide Synthesis," Springer Verlag, 1984; Bodanzsky, "Practice of Peptide Synthesis," Springer Verlag, 1984).

30 Accordingly, starting materials useful for preparing compounds of the invention and intermediates thereof are commercially available or can be prepared by well-known synthetic methods. Other methods for synthesis of the prodrugs described herein are either described in the art or will be readily apparent to the skilled artisan in view of the references provided above and may be used to synthesize the compounds of the invention.


Accordingly, the methods presented in the Schemes herein are illustrative rather than comprehensive.

In any of the Schemes below, after the amino group of a GABA analog has been functionalized with a promoiety or other protecting group, the carboxylic acid group may be converted to an ester or thioester by many synthetic methods, which are well-known to the skilled artisan. In one preferred embodiment, GABA analogs may be reacted with an alcohol or thiol in the presence of a coupling reagent (*e.g.*, carbodiimide and dimethylaminopyridine) to provide the ester. In another preferred embodiment, GABA analogs may be reacted with an alkyl halide in the presence of base to yield the ester. Other methods for converting GABA analogs to esters or thioesters are well within the purview of the skilled artisan in view of the references provided herein.


Scheme 1



(6)




(7)




As illustrated above in Scheme 1, carboxylic acids can be directly coupled to the terminal amino (or hydroxyl) group of GABA analog derivatives (6) to provide adducts (7). Reagents for effecting this reaction are well known to the skilled artisan and include, but are not limited to, carbodiimides, aminium salts, phosphonium salts, *etc.* Alternatively, reaction of carboxylic acid derivatives such as acyl chlorides, symmetrical anhydrides or mixed anhydrides with GABA analogs (6) in the presence of base (*e.g.*, hydroxide, tertiary amines, *etc.*) may be used to synthesize (7).

Scheme 2



As illustrated in Scheme 2, GABA analog derivatives (6) may be converted to carbamates (8) by treatment with various carbonic acid derivatives in the presence of base (e.g., hydroxide, tertiary amines, etc.). Alternatively, the well-known addition of alcohols to isocyanates (9) or (10) may be used to synthesize (8).

Scheme 3




As illustrated in Scheme 3, GABA analog derivatives (6) may be converted to thioamides (11) by reaction with thioacids in the presence of coupling agents. Reagents for effecting this reaction are well known to the skilled artisan and include, but are not limited to, carbodiimides, aminium salts, phosphonium salts, etc. Alternatively, reaction of thio acid derivatives such as thioacyl chlorides, symmetrical anhydrides or mixed anhydrides with (6) in the presence of base (e.g., hydroxide, tertiary amines, etc.) may be used to

synthesize thioamides (11). In yet another method, amides (7) may be converted to thioamides (11) by heating in the presence of phosphorus pentasulfide (when $n = 0$).

Thiocarbamates (12) and (13) may be synthesized from the reaction of the corresponding thiocarbonate derivatives (*i.e.*, $P = O$, $Q = S$ and $P = S$, $Q = O$, respectively) where W is chloride, imidazolyl or 4-nitrophenoxy with GABA analog derivatives (6) in the presence of base. Thiocarbamate (13) may also be formed by reaction of a thiol with isocyanates (9) or (10). Dithiocarbamate (14) ($P = S$, $Q = S$) may be made by reaction of GABA analog derivatives (6) with the dithiocarbonate derivative (*i.e.*, P and $Q = S$) where W is chloride, imidazolyl or 4-nitrophenoxy in the presence of base (see Scheme 4).



10

Scheme 4



One method for synthesis of compounds of Formula (IV) is illustrated in Scheme 5.


Scheme 5



Chloroformate (15) is treated with an aromatic leaving group such as *p*-nitrophenol in the presence of base to provide *p*-nitrophenylcarbonate (16). Halide interchange provides iodide (17), which is reacted with a metal or tetraalkylammonium salt of a carboxylic acid to afford compound (18). Treatment of (18) with GABA analog derivative (19), optionally in the presence of trimethylsilyl chloride, affords a compound of Formula (IV). Methods for making related acyloxyalkyl carbamate compounds have been described in the art (Alexander, United States Patent No. 4,760,057; Alexander, United States Patent No. 4,916,230; Alexander, United States Patent No. 5,466,811; Alexander, United States Patent No. 5,684,018).

Alternatively compounds of Formula (IV) can be prepared from carbonate (18) in a stepwise fashion as illustrated in Scheme 6. Here reaction of (18) with an α -amino acid (20), optionally protected as an ester, affords intermediate (21) which upon deprotection (if necessary) provides compound (22), which is then coupled to GABA analog (23) using standard peptide coupling reagents well known in the art.


Scheme 6



Another method for synthesis of compounds of Formula (IV) proceeds via carbonylation of GABA analog derivative (19) to an intermediate carbamic acid species, which is captured by an *in situ* alkylation reaction in an adaptation of the methods disclosed in the art (Butcher, *Synlett*, 1994, 825-6; Ferres *et al.*, U.S. Patent 4,036,829). Carbon dioxide gas is bubbled into a solution containing (19) and a base (*e.g.*, Cs₂CO₃, Ag₂CO₃ or AgO) in a solvent such as DMF or NMP. The activated halide is added, optionally in the

presence of iodide ion as a catalyst, and the carbonylation continued until the reaction is completed. This method is illustrated in Scheme 7 for the preparation of compounds of Formula (IV) from halide (24).

Scheme 7




X = Cl, Br

5

Alternatively compounds of Formula (IV) can be prepared in a stepwise fashion as illustrated in Scheme 8. Carbonylation and alkylation of carboxyl protected α -amino acid (20) provides intermediate (21), which upon deprotection is coupled to GABA analog (23) as previously described in Scheme 6.

10

Scheme 8




Yet another method for synthesis of compounds of Formula (IV) relies upon oxidation of ketocarbamate derivatives of GABA analogs (Gallop *et al.*, co-pending United States Patent Application Serial No. _____ entitled "Methods for Synthesis of Prodrugs from 1-Acyl-Alkyl Derivatives and Compositions Thereof"). As illustrated in Scheme 9, oxidation of ketocarbamate (25) affords compounds of Formula (IV). Preferred solvents include, but are not limited to, *t*-butanol, diethylether, acetic acid, hexane, dichloroethane, dichloromethane, ethyl acetate, acetonitrile, methanol, chloroform and water. Generally, the oxidant may be an organism (*e.g.*, yeast or bacteria), or a chemical reagent (*e.g.*, an enzyme or peroxide). Preferred oxidants include those, which have been successfully used in Baeyer-Villager oxidations of ketones to esters or lactones (Strukul, *Angew. Chem. Int. Ed.*, 1998, 37, 1198; Renz *et al.*, *Eur. J. Org. Chem.* 1999, 737; Beller *et al.*, in "Transition Metals in Organic Synthesis" Chapter 2, Wiley VCH; Stewart, *Current Organic Chemistry*, 1998, 2, 195; Kayser *et al.*, *Synlett*, 1999, 1, 153).

15

20


Scheme 9



Other compounds of this invention may be amenable to synthesis from the appropriate ketocarbamate derivative *via* this Baeyer-Villiger type oxidation, provided that they do not contain chemical functionality susceptible to decomposition or other transformation under conditions of the reaction.


Ketocarbamates (25) may be prepared from the corresponding α -hydroxyketone compounds (26) either directly, *via* reaction with isocyanate (9), or by first converting the α -hydroxyketone compound to a haloformate or activated carbonate intermediate (27) and subsequently reacting with compound (19), as illustrated in Scheme 10.

Scheme 10




Alternatively ketocarbamate (25) can be prepared in a stepwise manner via the α -amino acid carbamate (28) as illustrated in Scheme 11, following the coupling methodologies described above.

Scheme 11




Note that one method for preparation of isocyanate derivatives of GABA analogs (*i.e.*, compounds (9)) used in Scheme 10 above begins with the appropriate six-membered anhydride (29) as illustrated in Scheme 12. The anhydride ring is opened by reaction with an alcohol or thiol nucleophile to afford carboxylic acid (30). This compound is converted to the intermediate acyl azide in either a 2-step sequence (*i.e.*, first activation of the carboxyl group as a mixed anhydride, acyl halide or synthetic equivalent and then displacement with azide) or directly (*e.g.*, by treatment with Ph₂P(O)N₃). Curtius rearrangement of the acyl azide intermediate by thermolysis in an appropriate solvent (*e.g.*, toluene) at a temperature between 0 °C to 120 °C affords isocyanate (9). Optionally, the isocyanate is not isolated but rather is generated *in situ* and quenched by reaction with α-hydroxyketone (26) to afford the desired product (25).

Scheme 12



One method for synthesis of oxodioxolenylmethyl carbamate prodrugs (36) is disclosed in Scheme 13. Hydroxyketone (31) is treated with phosgene or carbonyldiimidazole in the presence of base to yield cyclic carbonate (32). Free radical bromination with N-bromosuccinimide and azoisobutyronitrile provides bromide (33), which is converted to alcohol (34). Alcohol (34) is transformed to dicarbonate (35) by reaction with 4-nitrophenyl chloroformate, which is then reacted with GABA analog derivatives (19) to provide prodrugs (36). Alternatively, reaction of compound (34) with isocyanate (9) provides compound (36), where n is 0.

Scheme 13




5

Prodrugs (41) may be synthesized by the method disclosed in Scheme 14.

Carboxylic acid (37) is coupled to alcohol (38) (e.g., dicyclohexylcarbodiimide and pyridine) to provide ester (39). Ester (39) is converted to activated carbonate (40) by reaction with 4-nitrophenyl chloroformate, which is then reacted with GABA analog derivative (19) to


10 provide prodrug (41).

Scheme 14




Enamine prodrugs such as (43) may be synthesized simply by reacting activated carbonyl compounds (42) with GABA analog derivatives (19) (where $R^{16} = H$), optionally in the presence of a secondary amine as catalyst, under dehydrating conditions as shown in Scheme 15.

Scheme 15




Compounds (III) may be synthesized by the route illustrated in Scheme 16. Reaction of GABA analog (23) with an α -activated ester derivative (44) provides amino ester (45). The amino group of (45) is blocked by acylation to yield (46) (e.g., using the methods described above) and the free acid is esterified under standard conditions to yield the diester (47). Dieckman condensation followed by decarboxylation yields ketone (48). Peroxy acid oxidation then provides lactone (III).

Scheme 16



Imine prodrugs (II) may be synthesized as depicted in Scheme 17 by treating ketones or ketone equivalents (49) with GABA analog derivatives (50) under dehydrating conditions.

Scheme 17



Phosphorus prodrugs may be synthesized by conventional methods known in the art. Similarly, prodrugs with S-N bond may be synthesized by using procedures described in the art.

5 4.4 Therapeutic Uses of the Compounds of the Invention

In accordance with the invention, a compound and/or composition of the invention is administered to a patient, preferably a human, suffering from epilepsy, depression, anxiety, psychosis, faintness attacks, hypokinesia, cranial disorders, neurodegenerative disorders, panic, pain (especially, neuropathic pain and muscular and skeletal pain), inflammatory
10 disease (*i.e.*, arthritis), insomnia, gastrointestinal disorders or ethanol withdrawal syndrome. Further, in certain embodiments, the compounds and/or compositions of the invention are administered to a patient, preferably a human, as a preventative measure against various diseases or disorders. Thus, the compounds and/or compositions of the invention may be administered as a preventative measure to a patient having a predisposition for epilepsy,
15 depression, anxiety, psychosis, faintness attacks, hypokinesia, cranial disorders, neurodegenerative disorders, panic, pain (especially, neuropathic pain and muscular and skeletal pain), inflammatory disease (*i.e.*, arthritis), insomnia, gastrointestinal disorders and ethanol withdrawal syndrome. Accordingly, the compounds and/or compositions of the invention may be used for the prevention of one disease or disorder and concurrently
20 treating another (*e.g.*, prevention of psychosis while treating gastrointestinal disorders; prevention of neuropathic pain while treating ethanol withdrawal syndrome).

The suitability of the compounds and/or compositions of the invention in treating epilepsy, depression, anxiety, psychosis, faintness attacks, hypokinesia, cranial disorders, neurodegenerative disorders, panic, pain (especially neuropathic pain and muscular and
25 skeletal pain), inflammatory disease (*i.e.*, arthritis), insomnia, gastrointestinal disorders and ethanol withdrawal syndrome may be determined by methods described in the art (See, *e.g.*, Satzinger *et al.*, United States Patent No. 4,024,175; Satzinger *et al.*, United States Patent No. 4,087,544; Woodruff, United States Patent No. 5,084,169; Silverman *et al.*, United States Patent No. 5,563,175; Singh, United States Patent No. 6,001,876; Horwell *et al.*,
30 United States Patent No. 6,020,370; Silverman *et al.*, United States Patent No. 6,028,214; Horwell *et al.*, United States Patent No. 6,103,932; Silverman *et al.*, United States Patent No. 6,117,906; Silverman, International Publication No. WO 92/09560; Silverman *et al.*, International Publication No. WO 93/23383; Horwell *et al.*, International Publication No. WO 97/29101, Horwell *et al.*, International Publication No. WO 97/33858; Horwell *et al.*,

International Publication No. WO 97/33859; Bryans *et al.*, International Publication No. WO 98/17627; Guglietta *et al.*, International Publication No. WO 99/08671; Bryans *et al.*, International Publication No. WO 99/21824; Bryans *et al.*, International Publication No. WO 99/31057; Magnus-Miller *et al.*, International Publication No. WO 99/37296; Bryans *et al.*, International Publication No. WO 99/31075; Bryans *et al.*, International Publication No. WO 99/61424; Pande, International Publication No. WO 00/23067; Bryans, International Publication No. WO 00/31020; Bryans *et al.*, International Publication No. WO 00/50027; and Bryans *et al.*, International Publication No. WO 02/00209). The compounds and/or compositions of the invention may be used to treat or prevent epilepsy, depression, anxiety, psychosis, faintness attacks, hypokinesia, cranial disorders, neurodegenerative disorders, panic, pain (especially neuropathic pain and muscular and skeletal pain), inflammatory disease (*i.e.*, arthritis), insomnia, gastrointestinal disorders and ethanol withdrawal syndrome by procedures described in the art (see references above). Thus, it is well with the capability of those of skill in the art to assay and use the compounds and/or compositions of the invention to treat or prevent epilepsy, depression, anxiety, psychosis, faintness attacks, hypokinesia, cranial disorders, neurodegenerative disorders, panic, pain (especially, neuropathic pain and muscular and skeletal pain), inflammatory disease (*i.e.*, arthritis), insomnia, gastrointestinal disorders and ethanol withdrawal syndrome.

20 4.5 Therapeutic/Prophylactic Administration

The compounds and/or compositions of the invention may be advantageously used in human medicine. As previously described in Section 4.4 above, compounds and/or compositions of the invention are useful for the treatment or prevention of epilepsy, depression, anxiety, psychosis, faintness attacks, hypokinesia, cranial disorders, neurodegenerative disorders, panic, pain (especially, neuropathic pain and muscular and skeletal pain), inflammatory disease (*i.e.*, arthritis), insomnia, gastrointestinal disorders or ethanol withdrawal syndrome.

When used to treat or prevent the above disease or disorders compounds and/or compositions of the invention may be administered or applied singly, in combination with other agents. The compounds and/or compositions of the invention may also be administered or applied singly, in combination with other pharmaceutically active agents, including other compounds of the invention.

The current invention provides methods of treatment and prophylaxis by administration to a patient of a therapeutically effective amount of a composition or

compound of the invention. The patient may be an animal, is more preferably a mammal, and most preferably a human.

The present compounds and/or compositions of the invention, which comprise one or more compounds of the invention, are preferably administered orally. The compounds and/or compositions of the invention may also be administered by any other convenient
5 route, for example, by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral mucosa, rectal and intestinal mucosa, *etc.*). Administration can be systemic or local. Various delivery systems are known, (*e.g.*, encapsulation in liposomes, microparticles, microcapsules, capsules, *etc.*) that can be used
10 to administer a compound and/or composition of the invention. Methods of administration include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, oral, sublingual, intranasal, intracerebral, intravaginal, transdermal, rectally, by inhalation, or topically, particularly to the ears, nose, eyes, or skin.

In particularly preferred embodiments, the compounds and/or compositions of the
15 invention can be delivered *via* sustained release systems, preferably oral sustained release systems. In one embodiment, a pump may be used (*see* Langer, *supra*; Sefton, 1987, *CRC Crit Ref Biomed Eng.* 14:201; Saudek *et al.*, 1989, *N. Engl. J Med.* 321:574).

In another embodiment, polymeric materials can be used (*see* "Medical Applications of Controlled Release," Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974);
20 "Controlled Drug Bioavailability," Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, 1983, *J Macromol. Sci. Rev. Macromol Chem.* 23:61; *see also* Levy *et al.*, 1985, *Science* 228: 190; During *et al.*, 1989, *Ann. Neurol.* 25:351; Howard *et al.*, 1989, *J. Neurosurg.* 71:105). In a preferred embodiment, polymeric materials are used for oral sustained release delivery. Preferred
25 polymers include sodium carboxymethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose and hydroxyethylcellulose (most preferred, hydroxypropylmethylcellulose). Other preferred cellulose ethers have been described (Alderman, *Int. J. Pharm. Tech. & Prod. Mfr.*, 1984, 5(3) 1-9). Factors affecting drug release are well known to the skilled artisan and have been described in the art (Bamba *et*
30 *al.*, *Int. J. Pharm.*, 1979, 2, 307).

In another embodiment, enteric-coated preparations can be used for oral sustained release administration. Preferred coating materials include polymers with a pH-dependent solubility (*i.e.*, pH-controlled release), polymers with a slow or pH-dependent rate of swelling, dissolution or erosion (*i.e.*, time-controlled release), polymers that are degraded by

enzymes (*i.e.*, enzyme-controlled release) and polymers that form firm layers that are destroyed by an increase in pressure (*i.e.*, pressure-controlled release).

In still another embodiment, osmotic delivery systems are used for oral sustained release administration (Verma *et al.*, *Drug Dev. Ind. Pharm.*, 2000, 26:695-708). In a preferred embodiment, OROSTM osmotic devices are used for oral sustained release delivery devices (Theeuwes *et al.*, United States Patent No. 3,845,770; Theeuwes *et al.*, United States Patent No. 3,916,899).

In yet another embodiment, a controlled-release system can be placed in proximity of the target of the compounds and/or composition of the invention, thus requiring only a fraction of the systemic dose (*see, e.g.*, Goodson, in "Medical Applications of Controlled Release," *supra*, vol. 2, pp. 115-138 (1984)). Other controlled-release systems discussed in Langer, 1990, *Science* 249:1527-1533 may also be used.

The compounds and/or compositions of the invention preferably provide GABA analogs (*e.g.*, gabapentin and pregablin) upon *in vivo* administration to a patient. While not wishing to bound by theory, the promoiety or promoieties of the compounds and/or compositions of the invention may be cleaved either chemically and/or enzymatically. One or more enzymes present in the stomach, intestinal lumen, intestinal tissue, blood, liver, brain or any other suitable tissue of a mammal may enzymatically cleave the promoiety or promoieties of the compounds and/or compositions of the invention. The mechanism of cleavage is not important to the current invention. Preferably, GABA analogs formed by cleavage of prodrugs from the compounds of the invention do not contain substantial quantities of lactam contaminant (preferably, less than 0.5 % by weight, more preferably, less than 0.2 % by weight, most preferably less than 0.1 % by weight). The extent of release of lactam contaminant from the prodrugs of this invention may be assessed using the standard *in vitro* analytical methods.

While not wishing to bound by theory, the promoiety or promoieties of the compounds and/or compositions of the invention may be cleaved prior to absorption by the gastrointestinal tract (*e.g.*, within the stomach or intestinal lumen) and/or after absorption by the gastrointestinal tract (*e.g.*, in intestinal tissue, blood, liver or other suitable tissue of a mammal). If the promoiety or promoieties of the compounds of the invention are cleaved prior to absorption by the gastrointestinal tract, the resulting GABA analogs may be absorbed into the systemic circulation conventionally (*e.g. via* the large neutral amino acid transporter located in the small intestine). If the promoiety or promoieties of the compounds of the invention are cleaved after absorption by the gastrointestinal tract, these

GABA analog prodrugs may have the opportunity to be absorbed into the systemic circulation either by passive diffusion, active transport or by both passive and active processes.

If the promoiety or promoieties of the compounds of the invention are cleaved after absorption by the gastrointestinal tract, these GABA analog prodrugs may have the opportunity to be absorbed into the systemic circulation from the large intestine. In this situation, the compounds and/or compositions of the invention are preferably administered as sustained release systems. In a preferred embodiment, the compounds and/or compositions of the invention are delivered by oral sustained release administration. Preferably, in this embodiment, the compounds and/or compositions of the invention are administered twice per day (more preferably, once per day).

4.6 Compositions of the Invention

The present compositions contain a therapeutically effective amount of one or more compounds of the invention, preferably in purified form, together with a suitable amount of a pharmaceutically acceptable vehicle, to provide the form for proper administration to a patient. When administered to a patient, the compounds of the invention and pharmaceutically acceptable vehicles are preferably sterile. Water is a preferred vehicle when a compound of the invention is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid vehicles, particularly for injectable solutions. Suitable pharmaceutical vehicles also include excipients such as starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The present compositions, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. In addition, auxiliary, stabilizing, thickening, lubricating and coloring agents may be used.

In one embodiment, the compositions of the invention are free of lactam side products formed by intramolecular cyclization. In a preferred embodiment, the compositions of the invention are stable to extended storage (preferably, greater than one year) without substantial lactam formation (preferably, less than 0.5% lactam by weight, more preferably, less than 0.2% lactam by weight, most preferably, less than 0.1% lactam by weight).

Pharmaceutical compositions comprising a compound of the invention may be manufactured by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Pharmaceutical compositions may be formulated in conventional manner using one or more
5 physiologically acceptable carriers, diluents, excipients or auxiliaries, which facilitate processing of compounds of the invention into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

The present compositions can take the form of solutions, suspensions, emulsion, tablets, pills, pellets, capsules, capsules containing liquids, powders, sustained-release
10 formulations, suppositories, emulsions, aerosols, sprays, suspensions, or any other form suitable for use. In one embodiment, the pharmaceutically acceptable vehicle is a capsule (see *e.g.*, Grosswald *et al.*, United States Patent No. 5,698,155). Other examples of suitable pharmaceutical vehicles have been described in the art (see Remington's Pharmaceutical Sciences, Philadelphia College of Pharmacy and Science, 17th Edition, 1985). Preferred
15 compositions of the invention are formulated for oral delivery, particularly for oral sustained release administration.

Compositions for oral delivery may be in the form of tablets, lozenges, aqueous or oily suspensions, granules, powders, emulsions, capsules, syrups, or elixirs, for example. Orally administered compositions may contain one or more optional agents, for example,
20 sweetening agents such as fructose, aspartame or saccharin, flavoring agents such as peppermint, oil of wintergreen, or cherry coloring agents and preserving agents, to provide a pharmaceutically palatable preparation. Moreover, where in tablet or pill form, the compositions may be coated to delay disintegration and absorption in the gastrointestinal tract, thereby providing a sustained action over an extended period of time. Selectively
25 permeable membranes surrounding an osmotically active driving compound are also suitable for orally administered compounds and compositions of the invention. In these later platforms, fluid from the environment surrounding the capsule is imbibed by the driving compound, which swells to displace the agent or agent composition through an aperture. These delivery platforms can provide an essentially zero order delivery profile as
30 opposed to the spiked profiles of immediate release formulations. A time delay material such as glycerol monostearate or glycerol stearate may also be used. Oral compositions can include standard vehicles such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, *etc.* Such vehicles are preferably of pharmaceutical grade.

For oral liquid preparations such as, for example, suspensions, elixirs and solutions, suitable carriers, excipients or diluents include water, saline, alkylene glycols (*e.g.*, propylene glycol), polyalkylene glycols (*e.g.*, polyethylene glycol) oils, alcohols, slightly acidic buffers between pH 4 and pH 6 (*e.g.*, acetate, citrate, ascorbate at between about 5 mM to about 50 mM), *etc.* Additionally, flavoring agents, preservatives, coloring agents, bile salts, acylcarnitines and the like may be added.

Compositions for administration via other routes may also be contemplated. For buccal administration, the compositions may take the form of tablets, lozenges, *etc.* formulated in conventional manner. Liquid drug formulations suitable for use with nebulizers and liquid spray devices and EHD aerosol devices will typically include a compound of the invention with a pharmaceutically acceptable vehicle. Preferably, the pharmaceutically acceptable vehicle is a liquid such as alcohol, water, polyethylene glycol or a perfluorocarbon. Optionally, another material may be added to alter the aerosol properties of the solution or suspension of compounds of the invention. Preferably, this material is liquid such as an alcohol, glycol, polyglycol or a fatty acid. Other methods of formulating liquid drug solutions or suspension suitable for use in aerosol devices are known to those of skill in the art (see, *e.g.*, Biesalski, United States Patent No. 5,112,598; Biesalski, United States Patent No. 5,556,611). A compound of the invention may also be formulated in rectal or vaginal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides. In addition to the formulations described previously, a compound of the invention may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, a compound of the invention may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

When a compound of the invention is acidic, it may be included in any of the above-described formulations as the free acid, a pharmaceutically acceptable salt, a solvate or hydrate. Pharmaceutically acceptable salts substantially retain the activity of the free acid, may be prepared by reaction with bases and tend to be more soluble in aqueous and other protic solvents than the corresponding free acid form.

4.7 Methods of Use And Doses

A compound of the invention, or compositions thereof, will generally be used in an amount effective to achieve the intended purpose. For use to treat or prevent diseases or disorders such as epilepsy, depression, anxiety, psychosis, faintness attacks, hypokinesia, cranial disorders, neurodegenerative disorders, panic, pain (especially neuropathic pain and muscular and skeletal pain), inflammatory disease (*i.e.*, arthritis), insomnia, gastrointestinal disorders or ethanol withdrawal syndrome the compounds of the invention or compositions thereof, are administered or applied in a therapeutically effective amount.

The amount of a compound of the invention that will be effective in the treatment of a particular disorder or condition disclosed herein will depend on the nature of the disorder or condition, and can be determined by standard clinical techniques known in the art as previously described. In addition, *in vitro* or *in vivo* assays may optionally be employed to help identify optimal dosage ranges. The amount of a compound of the invention administered will, of course, be dependent on, among other factors, the subject being treated, the weight of the subject, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

For example, the dosage may be delivered in a pharmaceutical composition by a single administration, by multiple applications or controlled release. In a preferred embodiment, the compounds of the invention are delivered by oral sustained release administration. Preferably, in this embodiment, the compounds of the invention are administered twice per day (more preferably, once per day). Dosing may be repeated intermittently, may be provided alone or in combination with other drugs and may continue as long as required for effective treatment of the disease state or disorder.

Suitable dosage ranges for oral administration are dependent on the potency of the parent GABA analog drug, but are generally about 0.001 mg to about 200 mg of a compound of the invention per kilogram body weight. When the GABA analog is gabapentin, typical daily doses of the parent drug in adult patients are 900 mg/day to 3600 mg/day and the dose of gabapentin prodrug may be adjusted to provide an equivalent molar quantity of gabapentin. Other GABA analogs may be more potent than gabapentin (*e.g.*, pregabalin), and lower doses may be appropriate for both the parent drug and any prodrug (measured on an equivalent molar basis). Dosage ranges may be readily determined by methods known to the skilled artisan.

The compounds of the invention are preferably assayed *in vitro* and *in vivo*, for the desired therapeutic or prophylactic activity, prior to use in humans. For example, *in vitro*

assays can be used to determine whether administration of a specific compound of the invention or a combination of compounds of the invention is preferred for reducing convulsion. The compounds of the invention may also be demonstrated to be effective and safe using animal model systems.

5 Preferably, a therapeutically effective dose of a compound of the invention described herein will provide therapeutic benefit without causing substantial toxicity. Toxicity of compounds of the invention may be determined using standard pharmaceutical procedures and may be readily ascertained by the skilled artisan. The dose ratio between toxic and therapeutic effect is the therapeutic index. A compound of the invention will
10 preferably exhibit particularly high therapeutic indices in treating disease and disorders. The dosage of a compound of the inventions described herein will preferably be within a range of circulating concentrations that include an effective dose with little or no toxicity.

4.8. Combination Therapy

15 In certain embodiments of the present invention, the compounds of the invention can be used in combination therapy with at least one other therapeutic agent. The compound of the invention and the therapeutic agent can act additively or, more preferably, synergistically. In a preferred embodiment, a composition comprising a compound of the invention is administered concurrently with the administration of another therapeutic agent,
20 which can be part of the same composition as the compound of the invention or a different composition. In another embodiment, a composition comprising a compound of the invention is administered prior or subsequent to administration of another therapeutic agent.

5. Examples

25 The invention is further defined by reference to the following examples, which describe in detail preparation of compounds and compositions of the invention and assays for using compounds and compositions of the invention. It will be apparent to those skilled in the art that many modifications, both to materials and methods, may be practiced without departing from the scope of the invention.

30 In the examples below, the following abbreviations have the following meanings. If an abbreviation is not defined, it has its generally accepted meaning.

35	AIBN	=	2, 2'-azobis(isobutyronitrile)
	Atm	=	atmosphere
	Boc	=	<i>tert</i> -butyloxycarbonyl
	Cbz	=	carbobenzyloxy

	CPM	=	counts per minute
	DCC	=	dicyclohexylcarbodiimide
	DMAP	=	4- <i>N,N</i> -dimethylaminopyridine
	DMEM	=	Dulbecco's minimum eagle medium
5	DMF	=	<i>N,N</i> -dimethylformamide
	DMSO	=	dimethylsulfoxide
	Fmoc	=	9-fluorenylmethyloxycarbonyl
	g	=	gram
	h	=	hour
10	HBSS	=	Hank's buffered saline solution
	L	=	liter
	LC/MS	=	liquid chromatography/mass spectroscopy
	M	=	molar
	min	=	minute
15	mL	=	milliliter
	mmol	=	millimoles
	NBS	=	N-bromosuccinimide
	NHS	=	N-hydroxysuccinimide
	PBS	=	phosphate buffered saline
20	THF	=	tetrahydrofuran
	TFA	=	trifluoroacetic acid
	TMS	=	trimethylsilyl
	μL	=	microliter
	μM	=	micromolar
25	v/v	=	volume to volume

EXAMPLE 1

1-[(α-Pivaloyloxymethoxy)carbonyl]aminomethyl}-1-Cyclohexane Acetic Acid (51)

30 Step A: Chloromethyl *p*-Nitrophenyl Carbonate (52)

p-Nitrophenol (100 g, 0.72 moles) was dissolved in anhydrous tetrahydrofuran (3 L) and stirred vigorously. To this solution was added chloromethyl chloroformate (70 mL, 0.79 moles) at room temperature followed by triethylamine (110 mL). After stirring for 1 hour, the reaction mixture was filtered and the filtrate was concentrated and then diluted

35 with ethyl acetate (1 L). The organic solution was washed with 10% potassium carbonate (3 x 500 mL) and 1N HCl (2 x 300 mL), brine (2 x 300 mL) and dried over anhydrous sodium sulfate. Removal of the solvent gave 157 g (95%) of the title compound (52) as a solid. The compound was unstable to LC-MS. ¹H NMR (CDCl₃, 400 MHz): 5.86 (s, 2H), 7.44 (d, *J* = 9 Hz, 2H), 8.33 (d, *J* = 9 Hz, 2H).

40

Step B: Iodomethyl *p*-Nitrophenyl Carbonate (53)

Chloromethyl *p*-nitrophenyl carbonate (52) (100 g, 0.43 moles), sodium iodide (228 g, 1.30 moles) and 50 g of dried molecular sieves (4Å) were added to 2 L of acetone under nitrogen with mechanical stirring. The resulting mixture was stirred at 40 °C for 5 hours (monitored by ¹H NMR). Upon completion, the solid materials were removed by filtration and the solvent was removed under reduced pressure. The residue was redissolved in dichloromethane (1 L) and washed twice with saturated aqueous sodium carbonate (300 mL) followed by water (300 mL). The organic layer was separated and dried over anhydrous sodium sulfate. Removal of solvent gave 123.6 g (89%) of the title compound (53) as a solid upon standing. The compound was found to be unstable to LC-MS. ¹H NMR (CDCl₃, 400 MHz): 6.06 (s, 2H), 7.42 (d, *J* = 9 Hz, 2H), 8.30 (d, *J* = 9 Hz, 2H). ¹³C NMR (CDCl₃, 100 MHz): 155.1, 151.0, 146.0, 125.8, 125.7, 121.9, 33.5.

Step C: Silver Pivalate (54)

Pivalic acid (50 g, 0.49 moles) was dissolved in acetonitrile (1.3 L) followed by addition of silver oxide (70 g, 0.29 moles) with vigorous stirring. Then, 660 mL of water was added under nitrogen. The resulting suspension was stirred at 70 °C in dark for 1 hour. After filtration through a pad of Celite, removal of the solvent gave 86 g (82%) of the title compound (54) as a pale white solid, which was used in the next reaction without further purification.

Other silver salts described in this application are prepared following similar procedures.

Step D: *p*-Nitrophenyl Pivaloyloxymethyl Carbonate (55)

To a solution of iodomethyl *p*-nitrophenyl carbonate (53) (62 g, 0.19 moles) in anhydrous toluene (1 L) was added silver pivalate (80 g, 0.38 moles). After stirring at 55 °C under nitrogen for 3 h, the reaction mixture was allowed to cool to room temperature and filtered through a pad of Celite. The filtrate was washed with 10% potassium carbonate (500 mL). Removal of the solvent yielded 43 g (75%) of the title compound (55) as a yellow oil. ¹H NMR (CDCl₃, 400 MHz): 1.25 (s, 9H), 5.88 (s, 2H), 7.40 (d, *J* = 9 Hz, 2H), 8.29 (d, *J* = 9 Hz, 2H). ¹³C NMR (CDCl₃, 100 MHz): 177.0, 155.3, 151.6, 145.8, 125.6, 121.9, 83.1, 39.1, 27.0.

Step E: 1-[[α -Pivaloyloxymethoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (51)

Gabapentin free base (24 g, 0.14 moles) was slurried in anhydrous dichloromethane (100 mL) and then treated with chlorotrimethylsilane (18.6 mL, 0.28 moles) and triethylamine (10 mL, 0.15 moles), respectively. The resulting suspension was warmed with stirring until complete dissolution of any solid was achieved. The above gabapentin solution was added *via* an equalizing addition funnel to a gently refluxed and mechanically stirred solution of *p*-nitrophenyl pivaloyloxymethyl carbonate (55) (20 g, 67 mmol) and triethylamine (10 mL, 0.15 moles) in dichloromethane (100 mL) under nitrogen. The resulting yellow solution was stirred for 1.5 hours. Upon completion (monitored by ninhydrin stain), the mixture was filtered and the filtrate was concentrated. The residue was dissolved in ethyl acetate (500 mL) and washed with 1N HCl (3 x 100 mL), brine (2 x 100 mL) and dried over anhydrous sodium sulfate. After removing the solvent, the crude product was dissolved in ethanol (300 mL) and then 1 g of 5% Pd/C was added. The resulting mixture was shaken under 50 psi hydrogen atmosphere for 15 minutes and then filtered through a pad of Celite. After concentration, the residue was dissolved in ethyl acetate, washed with 5% H₂SO₄ and dried over anhydrous sodium sulfate. After removing the solvent under reduced pressure, the residue was purified by chromatography on silica gel (4:1 hexanes :ethyl acetate) to afford 15 g (68%) of the title compound (51) as a solid. M.p.: 79-81 °C; ¹H NMR (CDCl₃, 400 MHz): 1.21 (s, 9H), 1.3-1.5 (m, 10H), 2.32 (s, 2H), 3.26 (s, 2H), 5.33 (m, 1H), 5.73 (s, 2H). ¹³C NMR (CDCl₃, 400 MHz): 21.7, 26.2, 27.3, 34.3, 38.2, 39.2, 80.6, 155.9, 176.8, 178.0. MS (ESI) m/z 328.36 (M-H)⁻, 330.32 (M+H)⁺, 352.33 (M+Na)⁺.

25

EXAMPLE 2

1-[[α -Acetoxyethoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (56)

Step A: 1-Chloroethyl-*p*-Nitrophenyl Carbonate (57)

To an ice cold reaction mixture containing *p*-nitrophenol (1.39 g, 10 mmol) and pyridine (0.81 g, 10 mmol) in dichloromethane (60 mL) was added 1-chloroethyl chloroformate (1.2 mL, 11 mmol). The mixture was stirred at 0 °C for 30 min and then at room temperature for 1 hour. After removing the solvent under reduced pressure, the residue was dissolved in ether, washed with water, 10% citric acid and water. The ether layer was dried over Na₂SO₄ and evaporated under reduced pressure to give 2.4 g (97%) of

the title compound (57) as an off-white solid. ¹H NMR (CDCl₃): 1.93 (d, 3H), 6.55 (q, 1H), 7.42 (d, 2H), 8.28 (d, 2H).

Step B: α-Acetoxyethyl-*p*-Nitrophenyl Carbonate (58)

5 A mixture of 1-chloroethyl-*p*-nitrophenyl carbonate (57) (0.5 g, 2 mmol) and mercuric acetate (1.5 g, 4.4 mmol) in acetic acid (15 mL) was stirred at room temperature for 24 hours. After removal of acetic acid under reduced pressure, the residue was dissolved in ether and washed with water, 0.5% (v/v) aqueous NaHCO₃, and water. The ether layer was dried over Na₂SO₄, and concentrated to dryness. Chromatography of the
10 resulting residue on silica gel, (hexanes: ethyl acetate (95:5)) gave 0.45 g (84%) of the title compound (58). ¹H NMR (CDCl₃, 400MHz): 1.55 (d, *J* = 5.6 Hz, 3H), 2.07 (s, 3H), 6.78 (q, *J* = 5.6 Hz, 1H), 7.36 (d, *J* = 9.6 Hz, 2H), 8.22 (d, *J* = 9.6 Hz, 2H).

Step C: 1-[(α-Acetoxyethoxy)carbonyl]aminomethyl}-1-Cyclohexane Acetic Acid (56)

15 To a mixture containing gabapentin (633 mg, 3.7 mmol) and triethylamine (1.03 mL, 7.4 mmol) in dichloromethane (20 mL) was added trimethylchlorosilane (0.93 mL, 7.4 mmol) and the mixture was stirred until a clear solution was formed. A solution containing α-acetoxyethyl-*p*-nitrophenyl carbonate (58) (1 g, 3.7 mmol) in dichloromethane (10 mL) was then added and the resulting mixture was stirred for 30 minutes. The reaction mixture
20 was washed with 10% citric acid (20 mL) and the organic layer was separated. The aqueous layer was further extracted with ether (3x10 mL) and the combined organic extracts were dried over MgSO₄. After filtration, the organic solvent was removed under reduced pressure. Chromatography of the resulting residue on silica gel, (hexanes: ethyl acetate (4:1)), gave 700 mg (63%) of the title compound (56). ¹H NMR (CDCl₃, 400MHz): 1.27-
25 1.60 (m, 10H), 1.55 (d, 3H), 2.08 (s, 3H), 2.38 (s, 2H), 3.25 (m, 2H), 5.31 (t, 1H), 6.81 (q, 1H). MS (ESI) *m/z* 302.22 (M+H)⁺. The acid form was quantitatively converted to the corresponding sodium salt by dissolution in water (5 mL), addition of an equimolar quantity of 0.5 N NaHCO₃, followed by lyophilization.

30

EXAMPLE 3

1-[(α-Benzoyloxybenzyloxy)carbonyl]aminomethyl}-1-Cyclohexane Acetic Acid (59)

Step A: *p*-Nitrophenyl α-Benzoylbenzylcarbonate (60)

To a solution of benzoin (2.0 g, 9.4 mmol) in 60 mL of CH_2Cl_2 was added DMAP (1.21 g, 9.9 mmol) and *p*-nitrophenyl-chloroformate (1.99 g, 9.9 mmol), respectively, at room temperature. After stirring for 3 hours at room temperature, the reaction was quenched with water and extracted with ethyl acetate/hexane (2x100 mL). The combined organic extracts were dried over anhydrous sodium sulfate. Removal of the solvent under reduced pressure afforded the title compound (60), which was used in the next reaction without purification.

Step B: 1-[(α -Benzoylbenzyloxy)carbonyl]aminomethyl}-1-Cyclohexane Acetic Acid
(61)

To a suspension of gabapentin (1.70 g, 9.9 mmol) in CH_2Cl_2 at 0°C was added triethylamine (2.76 mL, 19.8 mmol) and TMSCl (2.51 mL, 19.8 mmol). The reaction was then stirred for 30 minutes at room temperature. To this mixture was added compound (60) (prepared above in Step A) in CH_2Cl_2 and the resulting mixture was stirred at room temperature for 5 hours. The reaction mixture was diluted with dichloromethane, washed with brine and the organic phase was dried over Na_2SO_4 . After removing the solvent under reduced pressure, the residue was purified by chromatography on silica gel, eluting with 5% methanol in CH_2Cl_2 , to give 3.78 g (90% over two steps) of the title compound (61). ^1H NMR (CDCl_3 , 400MHz): δ 1.48-1.35 (m, 10H), 2.30 (s, 2H), 3.24 (d, $J = 7.2$ Hz, 2H), 5.58 (t, $J = 6.8$ Hz, 1H), 6.85 (s, 1H), 7.50-7.33 (m, 8H), 7.93 (d, $J = 7.2$ Hz, 2H).

Step C: 1-{[(α -Benzoyloxybenzyloxy)carbonyl]aminomethyl}-1-Cyclohexane Acetic Acid (59)

To a solution of 1- $\{[(\alpha\text{-benzoylbenzyloxy})\text{carbonyl}]\text{aminomethyl}\}$ -1-cyclohexane
25 acetic acid (**61**) (1.89 g, 4.6 mmol) in 40 mL of CH_2Cl_2 was added 77% *m*CPBA (2.07 g,
9.2 mmol) and NaHCO_3 (0.78 g, 9.2 mmol), respectively, at room temperature and the
resulting mixture was stirred at room temperature overnight. The reaction mixture was
acidified with 10% citric acid and extracted with CH_2Cl_2 . The organic extract was washed
with brine and dried over Na_2SO_4 . After removing the solvent under reduced pressure, the
30 residue was purified by reverse phase preparative HPLC (acetonitrile-water, 0.1% formic
acid) to afford 960 mg (49%) of the title compound (**59**). ^1H NMR (CDCl_3 , 400MHz): δ
1.58-1.35 (m, 10H), 2.34 (s, 2H), 3.26 (dd, $J = 6.8, 0.8$ Hz, 2H), 5.38 (t, $J = 6.8$ Hz, 1H),
7.46-7.26 (m, 5H), 7.63-7.55 (m, 3H), 7.89 (s, 1H), 8.08 (dd, $J = 8.8, 1.2$ Hz, 2H).

EXAMPLE 4

1-{[(α -Acetoxybenzyloxy)carbonyl]aminomethyl}-1-CyclohexaneAcetic Acid (62)

5 Following the procedure of Example 3, and substituting 1-hydroxy-1-phenyl-propan-2-one for benzoin, provided 300 mg of the title compound (62). ^1H NMR (CDCl_3 , 400MHz): δ 1.41 (m, 10H), 2.19 (s, 3H), 2.33 (s, 2H), 3.27 (dd, $J = 6.6, 1.6$ Hz, 2H), 5.36 (t, $J = 6.6$ Hz, 1H), 7.40 (m, 3H), 7.52 (m, 2H), 7.63 (s, 1H).

10

EXAMPLE 5

1-{[(α -Benzoyloxyethoxy)carbonyl]aminomethyl}-1-CyclohexaneAcetic Acid (63)

 Following the procedure of Example 3, and substituting 2-hydroxy-1-phenyl-1-propanone for benzoin, provided 5 mg of the title compound (63). ^1H NMR (CDCl_3 ,
15 400MHz): δ 1.44-1.36 (m, 10H), 1.62 (d, $J = 5.6$ Hz, 3H), 2.34 (s, 2H), 3.24 (d, $J = 6.8$ Hz, 2H), 5.28 (t, $J = 6.8$ Hz, 1H), 7.06 (q, $J = 5.6$ Hz, 1H), 7.44 (m, 2H), 7.56 (m, 1H), 8.03 (dd, $J = 8.4, 1.6$ Hz, 2H).

EXAMPLE 6

20

1-{[(1-Benzoyloxy-2-phenylethoxy)carbonyl]aminomethyl}-1-Cyclohexane Acetic Acid (64)Step A: 2-Phenyl-[1,3]-dithiane (65)

 To a solution of benzaldehyde (10.6 g, 100 mmol) and 1,3-propane dithiol in CH_2Cl_2
25 (150 mL) at room temperature was dropwise added $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (6.3 mL, 50 mmol) and the resulting mixture was stirred at room temperature for 2 hours. The reaction mixture was then diluted with CH_2Cl_2 , filtered and the filtrate washed with brine, saturated NaHCO_3 , brine and dried over Na_2SO_4 . The solvent was removed under reduced pressure to afford a white solid, which was recrystallized from a 1:1 mixture of ether and hexane to afford 17.0
30 g (87%) of the title compound (65) as white crystalline needles. ^1H NMR (CDCl_3 , 400MHz): δ 1.91 (m, 1H), 2.14 (m, 1H), 2.89 (m, 2H), 3.04 (m, 2H), 5.16 (s, 1H), 7.35-7.28 (m, 3H), 7.46 (m, 2H).

Step B: 2-Phenyl-1-(2-phenyl-[1,3]-dithian-2-yl)-ethanol (66)

To a solution of 2-phenyl-[1,3]-dithiane (65) (4.0 g, 20.4 mmol) in THF at -30°C was added a 1.6 M solution of n-butyllithium in THF (15.3 mL, 24.4 mmol). After stirring for 30 minutes at -30°C , a solution of phenylacetaldehyde (2.45 g, 20.4 mmol) in tetrahydrofuran was added dropwise at -30°C . The resulting reaction mixture was stirred for another hour at 0°C . The reaction was quenched with saturated NH_4Cl solution and extracted with ethyl acetate. The combined organic extracts were washed with saturated NH_4Cl solution, brine and dried over Na_2SO_4 . After filtrating and concentrating, the crude product was purified by flash chromatography on silica gel, (25% ethyl acetate in hexanes), to afford 2.63 g (71%) of the title compound (66). ^1H NMR (CDCl_3 , 400MHz): δ 1.97 (m, 2H), 2.23 (dd, $J = 4.0, 1.2$ Hz, 1H), 2.43 (dd, $J = 13.6, 10.2$ Hz, 1H), 2.77 (m, 4H), 3.02 (d, $J = 13.6$ Hz, 1H), 4.07 (m, 1H), 7.44-7.13 (m, 8H), 8.02 (dd, $J = 8.4, 1.4$ Hz, 2H).

Step C: 2-Hydroxy-1,3-diphenyl-propan-1-one (67)

To a solution of 2-phenyl-1-(2-phenyl-[1,3]-dithian-2-yl)-ethanol (66) (2.50 g, 7.9 mmol) in 100 mL of a 9:1 mixture of acetonitrile and water was added mercuric perchlorate hydrate (4.1 g, 10.3 mmol). The resulting mixture was stirred at room temperature for 5 minutes and thin layer chromatography indicated that the reaction was completed. The mixture was diluted with ethyl acetate, filtered through a pad of Celite and the filtrate was washed with saturated NaHCO_3 , brine and dried over Na_2SO_4 . The solvent was removed under reduced pressure and the crude product was purified by flash chromatography on silica gel, (20% ethyl acetate in hexanes) to afford 1.32 g (74%) of the title compound (67). ^1H NMR (CDCl_3 , 400MHz): δ 2.90 (dd, $J = 14.4, 7.0$ Hz, 1H), 3.20 (dd, $J = 14.4, 4.0$ Hz, 1H), 3.70 (d, $J = 6.8$ Hz, 1H), 5.35 (m, 1H), 7.28-7.11 (m, 5H), 7.53 (m, 2H), 7.65 (m, 1H), 7.93 (d, $J = 7.2$ Hz, 2H).

Step D: 1-[(1-Benzoyloxy-2-phenylethoxy)carbonyl]aminomethyl}-1-Cyclohexane Acetic Acid (64)

Following the procedure of Example 3, and substituting 2-hydroxy-1,3-diphenyl-propan-1-one for benzoin, provided 181 mg of the title compound (64). ^1H NMR (CDCl_3 , 400MHz): δ 1.45-1.29 (m, 10H), 2.24 (d, $J = 13.6$ Hz, 1H), 2.28 (d, $J = 13.6$ Hz, 1H), 3.22 (m, 4H), 5.26 (t, $J = 6.6$ Hz, 1H), 7.16 (t, $J = 5.6$ Hz, 1H), 7.33-7.25 (m, 5H), 7.40 (m, 2H), 7.57 (m, 1H), 8.02 (m, 2H).

EXAMPLE 7

**1-[[[(1-(3-Methylbutanoyloxy)-2-phenylethoxy)carbonyl]aminomethyl]-1-Cyclohexane
Acetic Acid (68)]**

Following the procedure of Example 6 and substituting 3-methylbutyraldehyde for benzaldehyde in Step A, provided 95 mg of the title compound (68). ¹H NMR (CDCl₃, 400MHz): δ 0.88-0.90 (m, 6H), 1.16-1.29 (m, 10H), 2.06 (m, 1H), 2.16 (m, 2H), 2.26 (m, 2H), 3.08 (d, *J* = 6.8 Hz, 2H), 3.19 (m, 2H), 5.22 (t, *J* = 6.8 Hz, 1H), 6.93 (t, *J* = 6 Hz, 1H), 7.31-7.23 (m, 5H).

EXAMPLE 8

1-[[[(α-Benzoyloxybutoxy)carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (69)]

Following the procedure of Example 6 and substituting butyraldehyde for phenylacetaldehyde in Step B, provided 240 mg of the title compound (69). ¹H NMR (CDCl₃, 400MHz): δ 0.99 (t, *J* = 7.6 Hz, 3H), 1.52-1.38 (m, 12H), 1.89 (m, 2H), 2.31 (s, 2H), 3.24 (m, 2H), 5.34 (t, *J* = 6.6 Hz, 1H), 6.70 (t, *J* = 5.6 Hz, 1H), 7.42 (m, 2H), 7.56 (m, 1H), 8.04 (m, 2H).

EXAMPLE 9

1-[[[(α-Acetoxybutoxy)carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (70)]

Following the procedure of Example 6, and substituting acetaldehyde for benzaldehyde in Step A and substituting butyraldehyde for phenylacetaldehyde in Step B respectively, provided 42 mg of the title compound (70). ¹H NMR (CD₃OD, 400MHz): δ 0.95 (m, 3H), 1.52-1.31 (m, 12H), 1.72 (m, 2H), 2.02 (s, 3H), 2.27 (s, 2H), 3.20 (s, 2H), 6.67 (t, *J* = 5.6 Hz, 1H).

EXAMPLE 10

**1-[[[(α-Butanoyloxybutoxy)carbonyl]aminomethyl]-1-Cyclohexane
Acetic Acid (71)]**

Following the procedure of Example 3, and substituting butyrolin for benzoin, provided 210 mg of the title compound (71). ¹H NMR (CDCl₃, 400 MHz); δ 0.93 (m, 6H), 1.37-1.76 (m, 16H), 2.30 (m, 4H), 3.23 (m, 2H), 5.25 (broad triplet, 1H), 6.73 (m, 1H). MS (ESI) *m/z* 356.45 (M-H)⁺.

EXAMPLE 11

1- $\{[(\alpha\text{-Propanoyloxyethoxy})\text{carbonyl}]\text{aminomethyl}\}$ -1-Cyclohexane Acetic Acid (72)**Step A: 1-Iodoethyl-*p*-Nitrophenyl Carbonate (73)**

5 A mixture of 1-chloroethyl-*p*-nitrophenyl carbonate (0.5 g, 2 mmol) and NaI (0.6 g, 4 mmol) in dry acetone was stirred for 3 hours at 40 °C. After filtration, the filtrate was concentrated under reduced pressure to afford 480 mg (72%) of the title compound (73), which was used in the next reaction without further purification.

Step B: α -Propanoyloxyethyl-*p*-Nitrophenyl Carbonate (74)

10 A mixture of 1-iodoethyl-*p*-nitrophenyl carbonate (73) (0.51 g, 1.5 mmol) and silver propionate (0.54 g, 3 mmol) in toluene (20 mL) was stirred at 50 °C for 24 hours. The reaction mixture was filtered to remove solids and the filtrate concentrated under reduced pressure. Chromatography of the resulting residue on silica gel, (20% CH₂Cl₂/hexanes and
15 then 40% CH₂Cl₂/hexanes), gave 0.39 g (92%) of the title compound (74). ¹H NMR (CDCl₃, 400MHz): 1.16 (t, *J* = 7.6 Hz, 3H), 1.61 (d, *J* = 5.6 Hz, 3H), 2.41 (q, *J* = 7.6 Hz, 2H), 6.84 (q, 1H, *J* = 5.6 Hz), 7.39 (d, *J* = 9.2 Hz, 2H), 8.28 (d, *J* = 9.2 Hz, 2H).

Step C: 1- $\{[(\alpha\text{-Propanoyloxyethoxy})\text{carbonyl}]\text{aminomethyl}\}$ -1-Cyclohexane Acetic Acid (72)

20 To a mixture of gabapentin (160 mg, 2.76 mmol) and triethylamine (0.77 mL, 5.5 mmol) in dichloromethane (30 mL) was added trimethylchlorosilane (0.71 mL, 5.5 mmol) and the resulting mixture was stirred until a clear solution was formed. To the above solution was added a solution of α -propanoyloxyethyl-*p*-nitrophenyl carbonate (74) (0.39g,
25 1.4 mmol) in dichloromethane (10 mL). After stirring for 30 minutes the reaction mixture was washed with 10% citric acid (20 mL) and the organic layer was separated. The aqueous layer was further extracted with ether (3 x 10 mL) and the combined organic extracts were dried over MgSO₄. After removing the solvent under reduced pressure, the residue was purified by reverse phase preparative HPLC (acetonitrile, water, 1% formic acid) to afford
30 190 mg (44%) of the title compound (72). ¹H NMR (CD₃OD, 400 MHz): 1.09 (t, *J* = 7.6 Hz, 3H), 1.36-1.54 (m, 10H), 1.44 (d, *J* = 5.6 Hz, 3H), 2.28 (s, 2H), 2.31 (q, *J* = 7.6 Hz, 2H), 3.22 (s, 2H), 6.67 (q, *J* = 5.6 Hz, 1H). MS (ESI) *m/z* 316.25 (M+H)⁺.

EXAMPLE 12

1-[(α -Butanoyloxyethoxy)carbonyl]aminomethyl}-1-Cyclohexane Acetic Acid (75)**Step A: α -Butanoyloxyethyl-*p*-Nitrophenyl Carbonate (76)**

5 A mixture of 1-iodoethyl-*p*-nitrophenyl carbonate (73) (1.5 g, 4.5 mmol) and silver butyrate (1.3 g, 6.7 mmol) in toluene (40 mL) was stirred at 90 °C in an oil bath for 24 h. The reaction mixture was filtered and the filtrate was concentrated under reduced pressure. Chromatography of the resulting residue on silica gel, (20% CH₂Cl₂/hexanes and then 40% CH₂Cl₂/hexanes), gave 0.46 g (36%) of the title compound (76). ¹H NMR (CDCl₃, 400
10 MHz): 0.95 (t, *J* = 7.6 Hz, 3H), 1.61 (d, *J* = 5.6 Hz, 3H), 1.67 (m, 2H), 2.41 (t, *J* = 7.6 Hz, 2H), 6.84 (q, 1H, *J* = 5.6 Hz), 7.39 (d, *J* = 9.2 Hz, 2H), 8.28 (d, *J* = 9.2 Hz, 2H). MS (ESI) *m/z* 298.28 (M+H)⁺.

Step B: 1-[(α -Butanoyloxyethoxy)carbonyl]aminomethyl}-1-Cyclohexane Acetic Acid (75)

15 To a mixture containing gabapentin (530 mg, 3.1 mmol) and triethylamine (0.89 mL, 6.4 mmol) in dichloromethane (30 mL) was added trimethylchlorosilane (0.83 mL, 6.4 mmol) and the resulting mixture was stirred until a clear solution was formed. To this solution was added a solution of α -butanoyloxyethyl-*p*-nitrophenyl carbonate (76) (0.46 g, 1.6 mmol) in dichloromethane (10 mL) and the resulting mixture was stirred for 30 min.
20 The reaction mixture was washed with 10% citric acid (20 mL) and the organic phase was separated. The aqueous layer was further extracted with ether (3 x 10 mL) and the combined organic phases were dried over MgSO₄, then concentrated *in vacuo*. The resulting residue was purified by reverse phase preparative HPLC (acetonitrile, water .1% formic acid) to afford 70 mg (21%) of the title compound (75). ¹H NMR (CD₃OD, 400MHz): 0.95 (t, *J* = 7.6 Hz, 3H), 1.32-1.58 (m, 10H), 1.42 (d, *J* = 5.6 Hz, 3H), 1.67 (m, 2H), 2.24 (s, 2H), 2.30 (t, *J* = 7.6 Hz, 2H), 3.24 (s, 2H), 6.74 (q, *J* = 5.6 Hz, 1H). MS (ESI) *m/z* 330.28 (M+H)⁺.
25

The acid form was quantitatively converted to the corresponding sodium salt by
30 dissolution in water (5 mL), addition of an equimolar quantity of 0.5 N NaHCO₃, followed by lyophilization.

EXAMPLE 13

1-[[α -Isobutanoyloxyethoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (77)

Following the procedure of Example 12, and substituting silver isobutyrate for silver butyrate, provided 70 mg (21%) of the title compound (77). ¹H NMR (CD₃OD, 400 MHz):
5 1.12 (d, *J* = 7.2 Hz, 3H), 1.14 (d, *J* = 7.2 Hz, 3H), 1.32-1.58 (m, 10H), 1.44 (d, *J* = 5.6 Hz, 3H), 2.28 (s, 2H), 2.56 (m, 1H), 3.25 (m, 2H), 6.73 (q, *J* = 5.6 Hz, 1H). MS (ESI) *m/z* 330.30 (M+H)⁺.

The acid form was quantitatively converted to the corresponding sodium salt by dissolution in water (5 mL), addition of an equimolar quantity of 0.5 N NaHCO₃, followed
10 by lyophilization.

EXAMPLE 14

1-[[α -Pivaloxyethoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (78)

Following the procedure of Example 12, and substituting silver pivalate for silver butyrate, provided 80 mg (36%) of the title compound (78). ¹H NMR (CDCl₃, 400MHz):
15 1.13 (s, 9H), 1.32-1.58 (m, 10H), 1.41 (d, *J* = 5.6 Hz, 3H), 2.27 (s, 2H), 3.25 (m, 2H), 5.41 (t, 1H), 6.73 (q, *J* = 5.6 Hz, 1H). MS (ESI) *m/z* 344.20 (M+H)⁺.

The acid form was quantitatively converted to the corresponding sodium salt by dissolution in water (5 mL), addition of an equimolar quantity of 0.5 N NaHCO₃, followed
20 by lyophilization.

EXAMPLE 15

1-[[α -Acetoxysisobutoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (79)

Following the procedure of Example 2, and substituting 1-chloro-2-methylpropyl chloroformate for 1-chloroethyl chloroformate, provided 212 mg (38%) of the title
25 compound (79). ¹H NMR (CD₃OD, 400 MHz): 0.99 (m, 6H), 1.32-1.58 (m, 10H), 1.88 (m, 1H), 2.08 (s, 3H), 2.38 (s, 2H), 3.25 (s, 2H), 6.52 (d, *J* = 4.4 Hz, 1H); MS (ESI) *m/z* 330.30 (M+H)⁺.

30

EXAMPLE 16

1-[[α -Pr panoyloxyisobutoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (80)

Following the procedure of Example 11, and substituting 1-chloro-2-methylpropyl-*p*-nitrophenyl carbonate for 1-chloroethyl-*p*-nitrophenyl carbonate, provided 190 mg (44%)

of the title compounds (80). ¹H NMR (CD₃OD, 400 MHz): 0.90 (d, *J* = 6.6 Hz, 3H), 0.91 (d, *J* = 6.6 Hz, 3H), 0.98 (t, *J* = 7.6 Hz, 3H), 1.32-1.58 (m, 10H), 1.83 (m, 1H), 2.18 (s, 2H), 2.28 (q, *J* = 7.6 Hz, 2H), 3.25 (s, 2H), 6.52 (d, *J* = 4.4 Hz, 1H). MS (ESI) *m/z* 344.34 (M+H)⁺.

5

EXAMPLE 17

1-(((α-Butanoyloxyisobutoxy)carbonyl)aminomethyl)-1-Cyclohexane Acetic Acid (81)

Following the procedure of Example 2 and substituting 1-chloro-2-methylpropyl chloroformate and mercuric butyrate for 1-chloroethyl chloroformate and mercuric acetate, respectively, provided 95 mg (36%) of the title compound (81). ¹H NMR (CD₃OD, 400 MHz): 1.12 (t, *J* = 7.6 Hz, 3H), 1.13 (d, *J* = 6.6 Hz, 3H), 1.14 (d, *J* = 6.6 Hz, 3H), 1.32-1.58 (m, 10H), 1.87 (m, 2H), 2.22 (m, 1H), 2.42 (s, 2H), 2.46 (t, *J* = 7.6 Hz, 2H), 3.44 (m, 2H), 6.78 (d, *J* = 4.8 Hz, 1H). MS (ESI) *m/z* 358.30 (M+H)⁺.

15

EXAMPLE 18

1-(((α-Isobutanoyloxyisobutoxy)carbonyl)aminomethyl)-1-Cyclohexane Acetic Acid (82)

Following the procedure of Example 2, and substituting 1-chloro-2-methylpropyl chloroformate and mercuric isobutyrate for 1-chloroethyl chloroformate and mercuric acetate, respectively, provided 95 mg (36%) of the title compound (82). ¹H NMR (CD₃OD, 400 MHz): 0.95 (d, *J* = 7.2 Hz, 3H), 0.97 (d, *J* = 7.2 Hz, 3H), 1.05 (d, *J* = 6.6 Hz, 3H), 1.06 (d, *J* = 6.6 Hz, 3H), 1.32-1.58 (m, 10H), 1.98 (m, 1H), 2.24 (s, 2H), 2.45 (m, 1H), 3.24 (m, 2H), 6.42 (d, *J* = 4.8 Hz, 1H). MS (ESI) *m/z* 358.27 (M+H)⁺.

25

EXAMPLE 19

1-(((α-Pivaloxyisobutoxy)carbonyl)aminomethyl)-1-Cyclohexane Acetic Acid (83)

Following the procedure of Example 12, and substituting 1-chloro-2-methylpropyl-*p*-nitrophenyl carbonate and silver pivalate for 1-chloroethyl-*p*-nitrophenyl carbonate and silver butyrate, respectively, provided 10 mg (9%) of the title compound (83). ¹H NMR (CD₃OD, 400 MHz): 0.98 (d, *J* = 6.6 Hz, 3H), 0.99 (d, *J* = 6.6 Hz, 3H), 1.19 (s, 9H), 1.32-1.58 (m, 10H), 2.08 (m, 1H), 2.28 (s, 2H), 3.21 (m, 2H), 6.49 (d, 1H); MS (ESI) *m/z* 372.31 (M+H)⁺.

30

EXAMPLE 20

1-[[α -Benzoyloxyisobutoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (84)

Following the procedure of Example 11, and substituting 1-chloro-2-methylpropyl-*p*-nitrophenyl carbonate and silver benzoate for 1-chloroethyl 1-*p*-nitrophenyl carbonate and silver propionate, respectively, provided 109 mg (40%) of the title compound (84). ¹H NMR (CD₃OD, 400 MHz): 1.18 (d, *J* = 7.2 Hz, 6H), 1.32-1.58 (m, 10H), 2.42 (m, 1H), 2.28 (s, 2H), 3.45 (s, 2H), 6.99 (d, *J* = 4.8 Hz, 1H), 7.76 (m, 2H), 7.92 (m, 1H), 8.26 (m, 2H). MS (ESI) *m/z* 392.22 (M+H)⁺.

EXAMPLE 21

1-[[α -Acetoxyisopropoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (85)**Step A: Isopropenyl-*p*-Nitrophenyl Carbonate (86)**

To a mixture of *p*-nitrophenol (5.76 g, 41.5 mmol) and isopropenyl chloroformate (5 g, 41.5 mmol) in dichloromethane (200 mL) at 0 °C was added a solution of pyridine (3.4 mL, 42 mmol) in dichloromethane (50 mL). The resulting mixture was stirred at 0 °C for 30 minutes and then at room temperature for 1 hour. After removing the solvent under reduced pressure, the residue was dissolved in ether and washed with water, 10% citric acid and water again. The ether layer was dried over Na₂SO₄ and evaporated under reduced pressure to give 8.7 g (94%) of the title compound (86) as an off-white solid. ¹H NMR (CDCl₃, 400 MHz): 2.05 (s, 3H), 4.81 (m, 1H), 4.95 (d, *J* = 2 Hz, 1H), 7.42 (d, *J* = 9.2 Hz, 2H), 8.28 (d, *J* = 9.2 Hz, 2H).

Step B: 2-Chloroisopropyl-*p*-Nitrophenyl Carbonate (87)

Isopropenyl-*p*-nitrophenyl carbonate (86) (8.7 g, 39 mmol) was dissolved in 4 M hydrogen chloride/dioxane in a sealed vessel. The mixture was stirred at room temperature for 16 hours. Removal of the solvent under reduced pressure gave 10 g (100%) of the title compound (87), which was used in the next reaction without further purification. ¹H NMR (CDCl₃, 400 MHz): 2.10 (s, 6H), 7.42 (d, 2H, *J* = 9.2 Hz), 8.28 (d, *J* = 9.2 Hz, 2H).

Step C: α -Acetoxyisopropyl-*p*-Nitrophenyl Carbonate (88)

A mixture of 2-chloroisopropyl-*p*-nitrophenyl carbonate (87) (0.5 g, 1.93 mmol) and mercuric acetate (1.0 g, 3.13 mmol) in dichloromethane (20 mL) was stirred at room temperature for 24 hours. The reaction mixture was filtered to remove solid and the filtrate

concentrated under reduced pressure. Chromatography of the resulting residue on silica gel, (20% CH₂Cl₂/hexanes and then 40% CH₂Cl₂/hexanes), afforded 227 mg (50%) of the title compound (**88**). ¹H NMR (CDCl₃, 400MHz): 1.90 (s, 6H), 2.07 (s, 3H), 7.28 (d, 2H, *J* = 9.2 Hz), 8.28 (d, *J* = 9.2 Hz, 2H).

5

Step D: 1-[[α -Acetoxyisopropoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (85)

To a mixture containing gabapentin (257 mg, 1.5 mmol) and triethylamine (0.46 mL, 3.3 mmol) in dichloromethane (30 mL) was added trimethylchlorosilane (0.38 mL, 3 mmol) and the mixture stirred until clear. A solution containing α -acetoxyisopropyl-*p*-nitrophenyl carbonate (**88**) (0.23 g, 0.8 mmol) in dichloromethane (10 mL) was added and stirred for 30 minutes. The reaction mixture was washed with brine (10 mL) and the organic layer was separated. The aqueous layer was further extracted with ether (3 x 10 mL) and the combined organic extracts were dried over MgSO₄ and then concentrated *in vacuo*. Chromatography of the resulting residue on silica gel, (hexane: ethyl acetate (4:1)), gave 40 mg (16%) of the title compound (**85**). ¹H NMR (CD₃OD, 400 MHz): 1.32-1.58 (m, 10H), 1.80 (s, 6H), 2.02 (s, 3H), 2.27 (s, 2H), 3.30 (s, 2H). MS (ESI) *m/z* 316.21 (M+H)⁺.

15

EXAMPLE 22

20 **1-[[α -Butanoyloxyisopropoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (89)**

Following the procedure of Example 21, and substituting mercuric butyrate for mercuric acetate, provided 5 mg (5%) of the title compound (**89**). ¹H NMR (CD₃OD, 400 MHz): 0.99 (t, *J* = 7.6 Hz, 3H), 1.32-1.58 (m, 10H), 1.60 (m, 2H), 1.85 (s, 6H), 2.22 (t, *J* = 7.6, 2H), 2.27 (s, 2H), 3.20 (s, 2H). MS (ESI) *m/z* 344.24 (M+H)⁺, 366.30 (M+Na)⁺.

25

EXAMPLE 23

1-[[α -Isobutanoyloxyisopropoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (90)

30 Following the procedure of Example 21, and substituting mercuric isobutyrate for mercuric acetate, provided 109 mg (43%) of the title compound (**90**). ¹H NMR (CD₃OD, 400MHz): 1.19 (d, *J* = 7.2 Hz, 6H), 1.32-1.58 (m, 10H), 1.82 (s, 6H), 2.38 (s, 2H), 3.25 (s, 2H). MS (ESI) 344.22 (M+H)⁺, 366.24 (M+Na)⁺.

EXAMPLE 24

1-[[α -Benzoyloxyisopropoxy]carbonyl]aminomethyl]-1-CyclohexaneAcetic Acid (91)

Following the procedure of Example 21, and substituting mercuric benzoate for mercuric acetate, provided 170 mg (58%) of the title compound (91). ^1H NMR (CDCl_3 , 400MHz): 1.32-1.58 (m, 10H), 1.95 (s, 6H), 2.30 (s, 2H), 3.20 (d, $J = 6.8$, 2H), 5.41 (t, $J = 6.8$ Hz, 1H), 7.40 (m, 2H), 7.52 (m, 1H), 7.98 (m, 2H). MS (ESI) m/z 400.29 ($\text{M} + \text{Na}$) $^+$.

EXAMPLE 25

1-[[α -Nicotinoyloxyisobutoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (92)Step A: 1-[[α -Chloroisobutoxy]carbonyl]aminomethyl-1-Cyclohexane Acetic Acid (93)

To a mixture containing gabapentin (1.71 g, 10 mmol) and triethylamine (3.06 mL, 22 mmol) in dichloromethane (150 mL) was added trimethylchlorosilane (1.4 mL, 11 mmol) and the resulting mixture was stirred until clear (about 20 min). A solution containing 1-chloro-2-methylpropylchloroformate (1.27 mL, 11 mmol) in dichloromethane (10 mL) was then added at 0 °C and stirred at room temperature for 60 min. The reaction mixture was washed with 10% citric acid (30 mL) and the organic layer separated. The aqueous layer was further extracted with ether (3x20 mL) and the combined organic phases were dried over MgSO_4 and then concentrated *in vacuo*. Chromatography of the residue on silica gel, eluting with hexane: ethyl acetate (1:4) gave 2.37g (77%) of the title compound. ^1H NMR (CDCl_3 , 400 MHz): δ 1.04 (d, $J = 6.4$ Hz, 3H), 1.06 (d, $J = 6.4$ Hz, 3H), 1.36-1.53 (m, 10H), 2.15 (m, 1H), 2.34 (s, 2H), 3.24 (m, 2H), 5.39 (t, 1H), 6.32 (d, $J = 5.6$ Hz, 1H). MS (ESI) m/z 306.34 ($\text{M} + \text{H}^+$).

Step B: 1-[[α -Nicotinoyloxyisobutoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (92)

A mixture of (93) (268 mg, 0.88 mmol), 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) (158 μL , 1.01 mmol), and nicotinic acid (637 mg, 5.2 mmol) in acetone was stirred at room temperature for 48 h. After filtration, the filtrate was concentrated *in vacuo* and the resulting residue was purified by reverse phase preparative HPLC to afford 50 mg (14%) of

the title compound. ^1H NMR (CD_3OD , 400 MHz): δ 1.07 (d, $J = 6.8$ Hz, 3H), 1.09 (d, $J = 6.8$ Hz, 3H), 1.32-1.58 (m, 10H), 2.19 (m, 1H), 2.26 (s, 2H), 3.23 (m, 2H), 6.78 (d, $J = 4.8$ Hz, 1H), 7.58 (m, 1H), 8.39 (d, $J = 6.4$ Hz, 1H), 8.76 (d, $J = 4.4$ Hz, 1H), 9.10 (s, 1H). MS (ESI) m/z 393.42 ($\text{M}+\text{H}^+$).

5

EXAMPLE 26

1-[[$(\alpha$ -2,2-Diethoxypropanoyloxyisobutoxy)carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (94)

10 Step A: Benzyl 1-[[$(\alpha$ -Chloroisobutoxy)carbonyl]aminomethyl]-1-Cyclohexane Acetate (95)

To a solution of (93) (1.02 g, 3.34 mmol) in dichloromethane was added 1,3-dicyclohexylcarbodiimide (758 mg, 3.67 mmol). After stirring at room temperature for 30 min, benzyl alcohol (380 μL , 3.67 mmol) and 4-(dimethylamino)pyridine (catalytic amount) were added. The resulting mixture was stirred at room temperature of 16 h. After filtration, the filtrate was washed with 10% citric acid, dried over Na_2SO_4 , and concentrated. Chromatography of the residue on silica gel, eluting with 10% ethyl acetate/hexane, gave 820 mg (62%) of the title compound. ^1H NMR (CDCl_3 , 400 MHz): δ 1.03 (d, $J = 6.4$ Hz, 3H), 1.05 (d, $J = 6.4$ Hz, 3H), 1.36-1.53 (m, 10H), 2.13 (m, 1H), 2.35 (s, 2H), 3.22 (m, 2H), 5.11 (s, 2H), 5.49 (t, 1H), 6.32 (d, $J = 4.8$ Hz, 1H), 7.34 (m, 5H). MS (ESI) m/z 396.24 ($\text{M}+\text{H}^+$).

20

Step B: Cesium 2,2-Diethoxypropionate (96)

To a stirred solution of 14 mL (0.2 mol) of pyruvic acid and 80 mL of triethylorthoformate at 10 $^\circ\text{C}$ was added 1 mL of concentrated sulfuric acid. The resulting mixture was stirred at 5-10 $^\circ\text{C}$ for 1 h and then diluted with 200 mL of dichloromethane. The organic solution was washed successively with water (3x80 mL) and saturated sodium chloride solution (80 mL) and then dried over anhydrous sodium sulfate. The mixture was filtered and then concentrated to give a quantitative yield of 2,2-diethoxypropionic acid as an oil. ^1H NMR (CDCl_3 , 400 MHz): δ 1.30 (t, 6H), 1.61 (s, 3H), 3.57 (q, 4H), 8.62 (s, 1H). The acid form was quantitatively converted to its cesium salt by dissolving the acid in water (25 mL) followed by treatment with an equimolar quantity of cesium carbonate, and then lyophilization. ^1H NMR (D_2O , 400 MHz): δ 0.98 (t, 6H), 1.28 (s, 3H), 3.22 (q, 2H), 3.47 (q, 2H).

30

Step C: Benzyl 1-{[(α -2,2-Diethoxypropanoyloxyisobutoxy)carbonyl]aminomethyl}-1-Cyclohexane Acetate (97)

A mixture of (95) (200mg, 0.51 mmol) and sodium iodide (114 mg, 0.76 mmol) in acetone was stirred at room temperature for 1 h. Cesium 2,2-diethoxypropionate (96) (300 mg, 1.02 mmol) and DMF (20 mL) were added and the resulting mixture was stirred at 40 °C for 18 h. After filtration, the filtrate was concentrated and the resulting residue was purified by silica gel flash column chromatography, eluting with 10% ethyl acetate/hexane to afford 100 mg (37%) of the title compound. MS (ESI) m/z 522.34 ($M+H^+$).

Step D: 1-{[(α -2,2-Diethoxypropanoyloxyisobutoxy)carbonyl]aminomethyl}-1-Cyclohexane Acetic Acid (94)

A mixture of (97) (200 mg, 0.38 mmol) and 5% Pd-C (catalytically amount) was stirred under hydrogen at room temperature for 16 h. After filtration, the filtrate was concentrated and the resulting residue was purified by reverse phase preparative HPLC to afford 98 mg (60%) of the title compound. 1H NMR ($CDCl_3$, 400 MHz): δ 0.97 (d, J = 6.8 Hz, 6H), 1.19 (t, J = 6.4 Hz, 3H), 1.21 (t, J = 6.4 Hz, 3H), 1.32-1.58 (m, 10H), 1.51 (s, 3H), 2.06 (m, 1H), 2.30 (s, 2H), 3.23 (m, 2H), 3.46 (m, 2H), 3.56 (m, 2H), 5.30 (t, 1H, NH), 6.59 (d, J = 4.8 Hz, 1H). MS (ESI) m/z 432.24 ($M+H^+$).

EXAMPLE 27

1-{[(α -(2-Amino-2-methylpropanoyl)oxyisobutoxy)carbonyl]aminomethyl}-1-Cyclohexane Acetic Acid (98)

Following the procedure of Example 26, and substituting 2-amino-2-methylpropionic acid for 2,2-diethoxypropionic acid, provided the title compound. 1H NMR ($CDCl_3$, 400 MHz): δ 0.97 (d, J = 6.8 Hz, 6H), 1.44 (s, 3H), 1.45 (s, 3H), 1.32-1.58 (m, 10H), 2.05 (m, 1H), 2.30 (s, 2H), 3.23 (m, 2H), 5.50 (t, 1H, NH), 6.58 (d, J = 4.8 Hz, 1H). MS (ESI) m/z 373.48 ($M+H^+$).

EXAMPLE 28

1-{[(α -Isobutanoyloxybutoxy)carbonyl]aminomethyl}-1-Cyclohexane Acetic Acid (99)

Step A: 2-Isopropyl-1,3-Dithiane (100)

To a mixture of isobutyraldehyde (9.1 mL, 100 mmol) and 1,3-propanedithiol (10 mL, 100 mmol) in dichloromethane at 0 °C was added boron trifluoride diethyl etherate (6.4 mL, 50 mmol). The resulting mixture was stirred at 0 °C for 30 min and at room temperature for 30 min. The reaction mixture was washed with brine, 5%NaHCO₃, and
5 brine again. The organic phase was separated and dried over Na₂SO₄, then concentrated to give 16 g (100%) of the title compound as a yellow liquid. This was carried to the next step without further purification. ¹H NMR (CDCl₃, 400 MHz): δ 1.057 (d, *J* = 7.2 Hz, 3H), 1.059 (d, *J* = 7.2 Hz, 3H), 1.80 (m, 1H), 1.97-2.08 (m, 2H), 2.82 (m, 4H), 4.00 (d, *J* = 5.2 Hz, 1H).

10 **Step B: 2-Isopropyl-2-(α-Hydroxybutyl)-1,3-Dithiane (101)**

To a solution of (100) (4 g, 24.7 mmol) in anhydrous tetrahydrofuran (50 mL) at –20°C was dropwise added *n*-butyl lithium (1.6M in hexane, 18.5 mL, 29.6 mmol). The stirred mixture was allowed to warm room temperature over 4 h and then cooled to –20 °C again. To this solution was added slowly a solution of *n*-butyraldehyde (2.7 mL, 29.6
15 mmol) in anhydrous tetrahydrofuran (10 mL). The resulting mixture was stirred for 16 h between –20 °C and room temperature. The reaction was quenched with saturated ammonium chloride solution and the mixture extracted with ethyl acetate. The organic layer was separated and dried over Na₂SO₄. After removing the solvent under reduced pressure, flash column chromatography of the residue on silica gel, eluting with 5% ethyl
20 acetate/hexane provided 5 g (85%) of the title compound as a yellow oil. ¹H NMR (CDCl₃, 400 MHz): δ 0.96 (t, *J* = 7.2 Hz, 3H), 1.11 (d, *J* = 6.8, Hz, 3H), 1.17 (d, *J* = 6.8 Hz, 3H), 1.42-1.52 (m, 2H), 1.76 (m, 1H), 1.87-1.95 (m, 2H), 2.04 (m, 2H), 2.62 (m, 4H), 2.94 (m, 2H), 4.03 (d, *J* = 5.2 Hz, 1H).

25 **Step C: 4-Hydroxy-2-Methylheptan-3-one (102)**

To a solution of (101) (5.0 g, 21.4 mmol) in acetonitrile (270 mL) was added under vigorous stirring a solution of Hg(ClO₄)₂ in methanol (30 mL). The resulting mixture was stirred at room temperature for 2 h. After filtration, the filtrate was carefully concentrated under reduced pressure without heating. Purification of the residue using silica gel flash
30 column chromatography (10% ethyl acetate/hexane) provided 2.8 g (91%) of the title compound as colorless liquid. ¹H NMR (CDCl₃, 400 MHz): δ 0.91 (t, *J* = 7.2 Hz, 3H), 1.09 (d, *J* = 7.2 Hz, 3H), 1.10 (d, *J* = 7.2 Hz, 3H), 1.35-1.46 (m, 4H), 1.75 (m, 1H), 2.80 (m, 1H), 3.45 (d, *J* = 5.2 Hz, 1H), 4.29 (m, 1H).

Step D: 2-Methylheptan-3-one-4-*p*-Nitrophenyl Carbonate (103)

To a mixture of (102) (1.1 g, 7.6 mmol), *p*-nitrophenyl chloroformate (1.84 g, 9.2 mmol) in anhydrous dichloromethane at 0 °C was added slowly a solution of 4-dimethylaminopyridine (1.12 g, 9.2 mmol) in dichloromethane. After stirring for 1 h at 0 °C and for 4 h at room temperature, the reaction was quenched with 10% citric acid. The organic phase was separated, dried over Na₂SO₄, and concentrated *in vacuo*. Flash column chromatography of the residue, eluting with 30% dichloromethane/hexane, provided 2 g (85%) of the title compound as an off-white solid. ¹H NMR (CDCl₃, 400 MHz): δ 0.99 (t, *J* = 7.6 Hz, 3H), 1.12 (d, *J* = 6.8 Hz, 3H), 1.18 (d, *J* = 6.8 Hz, 3H), 1.51 (m, 2H), 1.84 (m, 2H), 2.82 (m, 1H), 5.17 (m, 1H), 7.42 (d, *J* = 6.8 Hz, 2H), 8.25 (d, *J* = 6.8 Hz, 2H).

Step E: 1-[[α -Isobutanoylbutoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (104)

To a mixture containing gabapentin (820 mg, 4.8 mmol) and triethylamine (1.35 mL, 9.6 mmol) in dichloromethane (20 mL) was added trimethylchlorosilane (1.22 mL, 9.6 mmol) and the resulting mixture was stirred for 20 min. To this solution was added (103) (1 g, 3.2 mmol) in dichloromethane (10 mL) and the resulting mixture was stirred for 60 min. The reaction mixture was washed with 10% citric acid (20 mL) and the organic layer separated. The aqueous layer was further extracted with ether (3x10 mL) and the combined organic extracts were dried over MgSO₄ then concentrated *in vacuo*. Chromatography of the residue on silica gel, eluting with hexane: ethyl acetate (4:1) to remove *p*-nitrophenol, then further eluting with hexane: ethyl acetate (1:4) gave 780 mg (72%) of the title compound. ¹H NMR (CDCl₃, 400 MHz): δ 0.91 (t, *J* = 7.2 Hz, 3H), 1.04 (d, *J* = 6.8 Hz, 3H), 1.12 (d, *J* = 6.8 Hz, 3H), 1.36-1.53 (m, 12H), 1.74 (m, 2H), 2.33 (s, 2H), 2.78 (m, 1H), 3.22 (m, 2H), 5.11 (m, 1H), 5.48 (t, 1H, NH). MS (ESI) *m/z* 342.24 (M+H⁺).

Step F: 1-[[α -Isobutanoyloxybutoxy]carbonyl]aminomethyl]-1-Cyclohexane Acetic Acid (99)

To a solution of (104) (780 mg, 2.3 mmol) in dichloromethane (20 mL) was added *m*-chloroperoxybenzoic acid (1.03 g, 4.6 mmol) and NaHCO₃ (386 mg, 4.6 mmol). After stirring for 16 h at room temperature, another batch of *m*-chloroperoxybenzoic acid (791 mg, 4.6 mmol) and NaHCO₃ (386 mg, 4.6 mmol) was added. The resulting mixture was stirred for another 8 h and then treated with 10% citric acid. After filtration, the organic layer was separated, dried over Na₂SO₄, and concentrated. The residue was purified by

reverse phase preparative HPLC to afford 79 mg (11%) of the title compound. ¹H NMR (CDCl₃, 400 MHz): δ 0.94 (t, *J* = 7.2 Hz, 3H), 1.153 (d, *J* = 7.2 Hz, 3H), 1.150 (d, *J* = 7.2 Hz, 3H), 1.32-1.58 (m, 12H), 1.74 (m, 2H), 2.28 (s, 2H), 2.56 (m, 1H), 3.23 (m, 2H), 5.27 (t, *J* = 6.8 Hz, 1H, NH), 6.71 (t, *J* = 5.6 Hz, 1H). MS (ESI) *m/z* 358.30 (M+H⁺).

- 5 The above acid was quantitatively converted to the corresponding sodium salt by dissolving the acid in water (5 mL) followed by addition of an equimolar quantity of 0.5 N NaHCO₃ and lyophilization.

EXAMPLE 29

10 **Methyl 1-{[(α-Isobutanoyloxyisobutoxy)carbonyl]aminomethyl}-1-Cyclohexane**
Acetate (105)

Step A: **Methyl 1-{[(α-Chloroisobutoxy)carbonyl]aminomethyl}-1-Cyclohexane**
Acetate (106)

- 15 A mixture of (93) (1.0 g, 3.3 mmol), benzene (90 mL), and methanol (10 mL) was cooled to 0 °C. Trimethylsilyldiazomethane was added slowly at 0 °C until the yellow color persisted. The mixture was stirred at 0 °C for 30 min until the reaction was complete (monitored by TLC). After removing the solvent under reduced pressure, chromatography of the resulting residue on silica gel, eluting with 10% ethyl acetate/hexane gave 760 mg (72%) of the title compound. MS (ESI) *m/z* 320.24 (M+H⁺).

20

Step B: **Methyl 1-{[(α-Isobutanoyloxyisobutoxy)carbonyl]aminomethyl}-1-**
Cyclohexane Acetate (105)

- 25 A mixture of (106) (760 mg, 2.38 mmol), silver carbonate (394 mg, 1.4 mmol), and isobutyric acid (442 μL, 4.76 mmol) in chloroform was stirred at room temperature for 24 h. Another batch of silver carbonate (394 mg, 1.4 mmol) and isobutyric acid (442 μL, 4.76 mmol) was added, and the resulting mixture was stirred for another 24 h. After filtration, the filtrate was concentrated and the resulting residue purified by silica gel flash column chromatography, eluting with 10% ethyl acetate/hexane, to afford 560 mg (63%) of the title compound. ¹H NMR (CDCl₃, 400 MHz): δ 0.94 (d, *J* = 6.8 Hz, 3H), 0.96 (d, *J* = 6.8 Hz, 3H), 1.15 (d, *J* = 7.2 Hz, 3H), 1.17 (d, *J* = 7.2 Hz, 3H), 1.32-1.58 (m, 10H), 2.01 (m, 1H), 2.19 (s, 2H), 2.55 (m, 1H), 3.18 (m, 2H), 3.67 (s, 3H), 5.33 (t, 1H), 6.56 (d, *J* = 4.8 Hz, 1H). MS (ESI) *m/z* 372.38 (M+H⁺).
- 30

EXAMPLE 30

Methyl 1-[(α -Benzoyloxyisobutoxy)carbonyl]aminomethyl}-1-Cyclohexane Acetate
(107)

A mixture of 1-[(α -benzoyloxyisobutoxy)carbonyl]aminomethyl}-1-cyclohexane
5 acetic acid (**84**) (150 mg, 0.38 mmol), benzene (18 mL), and methanol (2 mL) was cooled to
0 °C. Trimethylsilyldiazomethane was added slowly at 0 °C until the yellow color persisted.
The mixture was stirred at 0 °C for 30 min until the reaction was complete (monitored by
TLC). After removing the solvent under reduced pressure, chromatography of the residue
on silica gel, eluting with 5% ethyl acetate/hexane gave 98 mg (64%) of the title compound.
10 ¹H NMR (CDCl₃, 400 MHz): δ 1.02 (d, J = 6.4 Hz, 3H), 1.03 (d, J = 6.4 Hz, 3H), 1.32-1.52
(m, 10H), 2.14 (m, 1H), 2.27 (s, 2H), 3.17 (m, 2H), 3.62 (s, 3H), 5.40 (t, 1H), 6.81 (d, J =
4.8 Hz, 1H), 7.40 (m, 2H), 7.54 (m, 1H), 8.12 (m, 2H). MS (ESI) m/z 406.29 (M+H⁺).

EXAMPLE 31

15 **1-[N-[(α -Isobutanoyloxyethoxy)carbonyl]-4-Bromophenylalaninyl] aminomethyl}-1-**
Cyclohexane Acetic Acid (108)

Step A: 1-[(4-Bromophenylalaninyl)aminomethyl]-1-Cyclohexane Acetate (109)

To a 40 mL vial was added an N-Boc-4-bromophenylalanine (1.72g, 5 mmol),
20 dicyclohexylcarbodiimide (1.24 g, 6 mmol), *N*-hydroxysuccinimide (0.7 g, 6 mmol), and
acetonitrile (20 mL). The reaction mixture was shaken at 25°C for 4 h. The precipitated
dicyclohexylurea was removed by filtration. To the filtrate was added an aqueous solution
(30 mL) of gabapentin hydrochloride (1.04 g, 6 mmol), and sodium hydroxide (0.4 g, 10
mmol). The reaction was stirred at 22-25 C for 16 h. The reaction mixture was diluted with
25 ethyl acetate (100 mL) and washed with 0.5 M aqueous citric acid (2x100 mL) and water
(2x100 mL). The organic phase was separated, dried (MgSO₄), filtered and concentrated
under reduced pressure. The residue was dissolved in trifluoroacetic acid (40 mL) and
allowed to stand at 22-25°C for 2 h. The solvent was removed under reduced pressure. The
residue was dissolved in water (4 mL) and filtered through a 0.25 μ m nylon membrane filter
30 prior to purification by preparative HPLC (Phenomenex 250x21.2 mm, 5 μ m LUNA C18
column, 100% water for 5 minutes, then 0-60% acetonitrile in water with 0.05% TFA over
20 minutes at 20 mL/min). The pure fractions were combined and the solvent was removed
under reduced pressure to afford 1.7 g (70%) of the title compound (**109**) as a white solid.
MS (ESI) m/z 397.02, 399.01 (M+H⁺).

Step B: 1-[[N-[(α -Isobutanoyloxyethoxy)carbonyl]-4-Bromophenylalaninyl]aminomethyl]-1-Cyclohexane Acetic Acid (108)

To a stirred suspension of (109) (200 mg, 0.51 mmol) in dichloromethane at 0 °C was added triethylamine (141 μ L, 1.01 mmol) and trimethylchlorosilane (129 mL, 1.01 mmol). The resulting mixture was stirred for 15 min at 0 °C, then a solution of α -isobutanoyloxyethyl-*p*-nitrophenyl carbonate (111) (144 mg, 0.51 mmol) in dichloromethane was added. The mixture was stirred at room temperature for 7 h (monitored by LC/MS) and then the reaction mixture was diluted with dichloromethane and acidified with citric acid. The organic layer was separated, washed with brine, and dried over Na₂SO₄. After filtration and concentration, the crude product was purified by preparative LC/MS to afford 92 mg of the title compound. ¹H-NMR (CD₃OD, 400MHz): δ 1.10 (m, 6H), 1.46-1.25 (m, 13 H), 2.20 (m, 2H), 2.48 (m, 1H), 2.84 (m, 1H), 3.06 (m, 1H), 3.17 (m, 1H), 4.36 (m, 1H), 6.67 (q, *J* = 5.6 Hz, 1H), 7.17 (d, *J* = 2.0, 8.0 Hz, 2H), 7.42 (dd, *J* = 2.0, 8.0 Hz, 2H).

EXAMPLE 32

3-[[(α -Isobutanoyloxyethoxy)carbonyl]aminomethyl]-5-Methylhexanoic Acid (110)

Step A: α -Isobutanoyloxyethyl-*p*-Nitrophenyl Carbonate (111)

A solution of 1-chloroethyl-*p*-nitrophenyl carbonate (57) (2.0 g, 8.14 mmol), and mercury isobutyrate (6.13 g, 16.29 mmol) in dichloromethane (10 mL) was stirred at 45 °C for 24 h. The reaction was then cooled to room temperature and diluted with hexane to precipitate mercury salts. The precipitate was filtered through a pad of Celite, and the filtrate was concentrated *in vacuo* to afford 2.5 g of crude product. Chromatography of the residue on silica gel, eluting with a gradient of 10% dichloromethane /hexane to 20% dichloromethane /hexane afforded 1.2 g (52%) of the title compound. ¹H-NMR (CDCl₃, 400MHz): δ 1.21-1.99 (m, 6H), 1.62 (d, *J* = 5.6 Hz, 3H), 2.61 (m, 1H), 6.84 (q, *J* = 5.6 Hz, 1H), 7.41 (dt, *J* = 6.8, 2.4 Hz, 2H), 8.29 (dt, *J* = 6.8, 2.4 Hz, 2H).

Step B: 3-[[(α -Isobutanoyloxyethoxy)carbonyl]aminomethyl]-5-Methylhexanoic Acid (110)

To a stirred suspension of pregabalin (2) (150 mg, 0.94 mmol) in anhydrous dichloromethane (10 mL) at 0 °C was added triethylamine (0.26 mL, 1.88 mmol) and

trimethylchlorosilane (0.24 mL, 1.88 mmol). After stirring for 15 min at 0 °C, a solution of α -isobutanoyloxyethyl-*p*-nitrophenyl carbonate (111) (267 mg, 0.94 mmol) in dichloromethane (3 mL) was added. The resulting mixture was stirred at room temperature for 1.5 h. The reaction mixture was acidified with citric acid and extracted with
5 dichloromethane. The combined organic extracts were washed with brine and dried over Na₂SO₄. After filtration and evaporation, the crude product was purified by silica gel chromatography, eluting first with dichloromethane to remove nitrophenol, then with 30% ethyl acetate in dichloromethane to afford 130 mg (48%) of the title compound as a mixture of two diastereomers. ¹H-NMR (CDCl₃, 400MHz): δ 0.90 (m, 6H), 1.70 (m, 8H), 1.46 (d, J = 5.6 Hz, 3H), 1.66 (1H, m), 2.15 (m, 1H), 2.33 (m, 2H), 2.53 (m, 1H), 3.12 (m, 1H), 3.29
10 (m, 1H), 5.08 (t, J = 6.0 Hz, 1H), 6.79 (m, 1H).

EXAMPLE 33

3-[(α -Isobutanoyloxyisobutoxy)carbonyl]aminomethyl]-5-Methyl-Hexanoic 15 Acid (112)

Step A: 1-Chloro-2-Methylpropyl-*p*-Nitrophenyl Carbonate (113)

To an ice cold reaction mixture containing *p*-nitrophenol (4.06 g, 29 mmol) and 1-chloro-2-methylpropyl chloroformate (5.0 g, 29 mmol) in dichloromethane (200 mL) was
20 added a solution of pyridine (2.78 mL, 32 mmol) in dichloromethane (50 mL). The mixture was stirred at 0 °C for 30 min and then at room temperature for 1 h. After evaporation of the solvent under reduced pressure, the residue was dissolved in ether and washed with water, 10% citric acid and water again. The ether layer was separated, dried over Na₂SO₄ and evaporated under reduced pressure to give 7.9 g (100%) of the title compound as an off-
25 white solid. ¹H NMR (CDCl₃, 400 MHz): δ 1.12 (d, J = 6.6 Hz, 3H), 1.13 (d, J = 6.6 Hz, 3H), 2.29 (m, 1H), 6.24 (d, J = 4.8 Hz, 1H), 7.42 (d, J = 9.2 Hz, 2H), 8.28 (d, J = 9.2 Hz, 2H).

Step B: α -Isobutanoyloxyisobutyl-*p*-Nitrophenyl Carbonate (114)

30 Following the procedure for preparation of (111), and substituting (113) for (57), provided the title compound in 15% yield with a 70% recovery of starting material. ¹H-NMR (CDCl₃, 400MHz): δ 1.07 (d, J = 6.8 Hz), 1.21 (m, 6H), 2.18 (m, 1H), 2.26 (m, 1H), 6.60 (d, J = 5.2 Hz, 1H), 7.42 (m, 2H), 8.28 (m, 2H).

Step C: 3-[[α -Isobutanoyloxyisobutoxy]carbonyl]aminomethyl]-5-Methyl-Hexanoic Acid (112)

Following the procedure of preparation for (110), and substituting (114) for (111), provided the title compound as a mixture of two diastereomers in 51% yield. ¹H-NMR (CDCl₃, 400MHz): δ 0.89 (m, 12H), 1.17 (m, 8H), 1.65 (m, 1H), 2.02 (m, 1H), 2.16 (m, 1H), 2.33 (m, 2H), 2.56 (m, 1H), 3.13 (m, 1H), 3.30 (m, 1H), 5.00 (m, 1H), 6.57-6.56 (m, 1H).

EXAMPLE 34

10 **3-[[α -Benzoyloxyisobutoxy]carbonyl]aminomethyl]-5-Methyl-Hexanoic Acid (115)**

Step A: α -Benzoyloxyisobutyl-*p*-Nitrophenyl Carbonate (116)

Following the procedure of preparation for (111), substituting (113) for (57) and mercury benzoate for mercury isobutyrate, provided the title compound in 11% yield with a 50% recovery of starting material. ¹H-NMR (CDCl₃, 400MHz): δ 1.15 (d, J = 3.2 Hz, 3H), 1.16 (d, J = 3.2 Hz, 3H), 2.30 (m, 1H), 6.87 (d, J = 4.4 Hz, 1H), 7.42 (dd, J = 7.2, 2.0 Hz, 2H), 7.48 (t, J = 7.6 Hz, 2H), 7.62 (t, J = 7.6 Hz, 1H), 8.09 (dd, J = 8.0, 1.0 Hz, 2H), 8.27 (dd, J = 7.2, 2.0 Hz, 2H).

20

Step B: 3-[[α -Benzoyloxyisobutoxy]carbonyl]aminomethyl]-5-Methyl-Hexanoic Acid (115)

Following the procedure of preparation for (110), and substituting (116) for (111), provided the title compound as a mixture of two diastereomers in 58% yield. ¹H-NMR (CDCl₃, 400MHz): δ 0.87(m, 6H), 1.05 (m, 6H), 1.16 (m, 2H), 1.64 (m, 1H), 2.17 (m, 2H), 2.32 (m, 2H), 3.12 (m, 1H), 3.29 (m, 1H), 5.01 (br s, 1H), 6.82 (m, 1H), 7.44 (m, 2H), 7.57 (m, 1H), 8.05 (m, 2H).

30

1-[[α -(5-Methyl-2-Oxo-1,3-Dioxol-4-en-4-yl)methoxy]carbonyl]aminomethyl]-1-Cycl hexane Acetic Acid (117)

Step A: Benzyl 2-Diazo-3-Ox -Butyric Acid (118)

To a solution of benzyl acetoacetate (5.0 g, 26.01 mmol) and 4-acetamido-benzenesulfonyl azide (6.25 g, 26.01 mmol) in acetonitrile (200 mL) at 0 °C was dropwise added triethylamine (10.9 mL, 78.03 mmol). The resulting mixture was stirred for 30 min at 0 °C and 4 h at room temperature. After concentrating under reduced pressure, the residue was triturated with 2:1 ethyl ether/petroleum ether (3x100 mL). The combined organic extract was filtered through a pad of Celite topped with silica gel. Removal of the solvent under reduced pressure afforded 4.74 g of the title compound as off-white crystals. ¹H-NMR (CDCl₃, 400MHz): δ 2.49 (s, 3H), 5.27 (s, 2H), 7.38 (m, 5H).

10 **Step B: Benzyl 2-Hydroxy-3-Oxo-Butyric Acid (119)**

A solution of the diazo compound (118) (4.74 g, 21.74 mmol) in THF (110 mL) and H₂O (50 mL) was heated under reflux with Rh₂(OAc)₂ (77 mg, 0.17 mmol) for 4 h and allowed to cool to room temperature. The mixture was concentrated *in vacuo* and the aqueous residue was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over Na₂SO₄, filtered and concentrated *in vacuo* to provide 4.5 g of crude product. ¹H-NMR (CDCl₃, 400MHz): δ 2.28 (s, 3H), 3.90 (s, 1H), 4.82 (s, 1H), 5.26 (m, 2H), 7.37 (m, 5H).

20 **Step C: 4-Benzyloxycarbonyl 5-Methyl-2-Oxo-1,3-Dioxol-4-ene (120)**

To a suspension of carbonyldiimidazole (6.88 g, 42.45 mmol) in THF (50 mL) at 0 °C was added a solution of alcohol (119) (4.50 g, 21.22 mmol) in dry THF (50 mL). The resulting mixture was stirred for 5 h at 0 °C, then overnight at room temperature. The mixture was concentrated *in vacuo* and the residue was partitioned with water and ethyl acetate/hexane. The organic layer was separated and washed with saturated NH₄Cl, brine, and dried over Na₂SO₄. After filtration and concentration, the crude product was purified by flash chromatography on silica gel, eluting with 20% ethyl acetate in hexane to afford 2.6 g of the title compound. ¹H-NMR (CDCl₃, 400MHz): δ 2.48 (s, 3H), 5.27 (s, 2H), 7.37 (br. s, 5H).

30 **Step D: 5-Methyl-2-Oxo-1,3-Dioxol-4-enyl-4-Carboxylic Acid (121)**

To a solution of compound (120) (2.6 g, 10.92 mmol) in 50 mL of ethanol was added 260 mg of Pd/C (5%) and the resulting mixture was stirred under a hydrogen atmosphere for 1 h. Filtration and removal of solvent under reduced pressure provided 1.62 g of the title compound. ¹H-NMR (CD₃OD, 400MHz): δ 2.41 (s, 3H).

Step E: 4-Hydroxymethyl-5-Methyl-2-Oxo-1,3-Dioxol-4-ene (122)

To a solution of the acid (121) (1.62 g, 11.10 mmol) and anhydrous DMF (112 μ L) in dry dichloromethane (50 mL) at 0 °C was dropwise added oxalyl chloride (6.1 mL of 2M solution, 12.2 mmol). After stirring for 30 min at 0 °C and 1 h at room temperature, the solvent was removed under reduced pressure. The residue was dissolved in anhydrous dichloromethane (65 mL) and cooled to -78 °C. To this solution was dropwise added a solution of Bu₄NBH₄ (3.14 g, 12.2 mmol, in 20 mL dichloromethane) over 10 min. After stirring for 1 h at -78 °C, the mixture was cautiously quenched with 0.1N HCl (30 mL) and allowed to warm to room temperature. The aqueous layer was separated and was extracted with EtOAc (3x50 mL) and the combined organic extract was washed with brine and dried over Na₂SO₄. After removing the solvent under reduced pressure, column chromatography on silica gel, eluting with 50% EtOAc in dichloromethane provided 767 mg of the title compound. ¹H-NMR (CD₃OD, 400MHz): δ 2.09 (s, 3H), 4.34 (s, 2H).

Step F: Benzyl 1-(((5-Methyl-2-Oxo-1,3-Dioxol-4-en-4-yl)methoxy)carbonyl)-aminomethyl}-1-Cyclohexane Acetate (123)

A suspension of the alcohol (122) (767 mg, 5.9 mmol) and benzyl 1-isocyanatomethyl-1-cyclohexane acetate (5.9 mmol) in toluene was refluxed overnight. After removing the solvent under reduced pressure, the residue was purified by flash column chromatography, eluting with 30% EtOAc in hexane to provide 510 mg of the title compound. ¹H-NMR (CD₃OD, 400MHz): δ 1.58-1.30 (m, 10H), 2.18 (s, 3H), 2.35 (s, 2H), 3.17 (d, J = 6.8 Hz, 2H), 4.80 (s, 2H), 5.11 (s, 2H), 5.44 (t, J = 6.8 Hz, 1H), 7.36 (m, 5H).

Step G: 1-(((5-Methyl-2-Oxo-1,3-Dioxol-4-en-4-yl)methoxy)carbonyl)-aminomethyl}-1-Cyclohexane Acetic Acid (117)

To a solution of compound (123) (510mg, 1.41 mmol) in ethanol (20 mL) was added 59 mg of Pd/C (5%) and the resulting mixture was stirred under a hydrogen atmosphere for 1 h. Filtration and removal of volatiles under reduced pressure provided the crude product, which was purified by preparative LC/MS to provide 105 mg of the title compound. ¹H-NMR (CD₃OD, 400MHz): δ 1.52-1.36 (m, 10H), 2.16 (s, 3H), 2.27 (s, 2H), 3.22 (s, 2H), 4.86 (s, 2H).

EXAMPLE 36

Piperidinium 1-[(1-Methyl-3-Oxo-But-1-enyl)aminomethyl]-1-Cyclohexane Acetate (124)

2,4-pentanedione (103 μ L, 1 mmol), gabapentin (171 mg, 1 mmol), and piperidine
5 (99 μ L, 1 mmol) were mixed in anhydrous methanol (10 mL). The resulting mixture was
heated under reflux for 4 h. Removal of the solvent under reduced pressure gave the title
compound with purity greater than 90%. ^1H NMR (CDCl_3 , 400MHz): δ 1.34-1.62 (m,
12H), 1.71 (m, 4H), 1.94 (s, 3H), 1.96 (s, 3H), 2.26 (s, 2H), 2.98 (m, 4H), 3.38 (d, J = 6 Hz,
2H), 4.90 (s, 1H), 5.20 (s, br, 2H), 8.64 (t, J = 6 Hz, 1H). MS (ESI) m/z 252.35 (M-H^+).
10

EXAMPLE 37

Piperidinium 1-[1-[(2-Oxo-Tetrahydrofuran-3-ylidene)ethyl]aminomethyl]-1-Cyclohexane Acetate (125)

2-Acetylbutyrolactone (108 μ L, 1 mmol), gabapentin (171 mg, 1 mmol), and
15 piperidine (99 μ L, 1 mmol) were mixed in anhydrous methanol (10 mL). After heating
under reflux for 6 h, the solvent was removed under reduced pressure to afford the title
compound with purity greater than 90%. ^1H NMR (CDCl_3 , 400MHz): δ 1.34-1.62 (m,
12H), 1.71 (m, 4H), 1.94 (s, 3H), 2.24 (s, 2H), 2.81 (t, J = 7.6 Hz, 2H), 2.99 (m, 4H), 3.31
(d, J = 6.4 Hz, 2H), 4.23 (t, J = 7.6 Hz, 2H), 5.17 (s, br, 2H), 8.64 (t, J = 6.4 Hz, 1H). MS
20 (ESI) m/z 280.34 (M-H^+).

EXAMPLE 38

Piperidinium 1-[(2-Carbomethoxy-Cyclopent-1-enyl)aminomethyl]-1-Cyclohexane Acetate (126)

25 Methyl 2-oxocyclopentanecarboxylate (124 μ L, 1 mmol), gabapentin (171 mg, 1
mmol), and piperidine (99 μ L, 1 mmol) were mixed in anhydrous methanol (10 mL). After
heating under reflux for 16 h, the solvent was removed under reduced pressure to afford the
title compound with purity greater than 90%. ^1H NMR (CDCl_3 , 400MHz): δ 1.29-1.60 (m,
12H), 1.72 (m, 4H), 1.79 (m, J = 7.6 Hz, 2H), 2.24 (s, 2H), 2.49 (t, J = 7.6 Hz, 2H), 2.55 (t,
30 J = 7.6 Hz, 2H), 2.99 (m, 4H), 3.24 (d, J = 6.8 Hz, 2H), 3.63 (s, 3H), 5.06 (s, br, 2H), 7.93
(s, br, 1H). MS (ESI) m/z 294.36 (M-H^+).

EXAMPLE 39

Piperidinium 1-[(1-Methyl-2-(Ethoxycarbonyl)-3-Ethoxy-3-Oxoprop-1-enyl)aminomethyl]-1-Cyclohexane Acetate (127)

Diethyl acetylmalonate (202 mg, 1 mmol), gabapentin (171 mg, 1 mmol), and
5 piperidine (99 μ L, 1 mmol) were mixed in anhydrous ethanol (10 mL). After heating under
reflux for 16 h, the solvent was removed under reduced pressure to give the title compound
with purity greater than 90%. ^1H NMR (CDCl_3 , 400MHz): δ 1.28 (t, $J = 7.2$ Hz, 6H), 1.38-
1.64 (m, 12H), 1.75 (m, 4H), 1.96 (s, 3H), 2.23 (s, 2H), 2.99 (m, 4H), 3.24 (d, $J = 5.2$ Hz,
2H), 4.20 (q, $J = 7.2$ Hz, 4H), 4.35 (s, br, 2H), 7.79 (t, $J = 5.2$ Hz, 1H). MS (ESI) m/z
10 354.38 (M-H $^+$).

EXAMPLE 40

1-[(α -(2-(2-Methyl-1,3-Dioxolan-2-yl)carboxyisobutoxy)carbonyl)aminomethyl]-1-Cyclohexane Acetic Acid (128)

15

Step A: 2-Methyl-1,3-Dioxolane-2-Carboxylic Acid (129)

To a stirred mixture containing ethyl pyruvate (11.1 mL, 0.1 mol) and ethylene
glycol (5.6 mL, 0.1 mol) in anhydrous dichloromethane (100 mL) at 0 $^\circ\text{C}$ was added boron
trifluoride dietherate (6.4 mL, 0.05 mol) and catalytic amount of acetic acid. The resulting
20 mixture was stirred at 40 $^\circ\text{C}$ for 16 h and then diluted with 100 mL of dichloromethane.
The organic solution was washed successively with saturated sodium chloride solution (2 x
80 mL). The organic layer was separated and the combined organic extracts were
concentrated. The residue was treated with 1N sodium hydroxide at room temperature.
After stirring at room temperature for 3 h (monitored by TLC), citric acid was added to
25 adjust the pH to 4. The product was extracted with dichloromethane, dried over Na_2SO_4
and concentrated to afford 5.1 g (38%) of the title compound (129) as a clear liquid. This
material was used in the next reaction without further purification. ^1H NMR (CDCl_3 , 400
MHz): δ 1.55 (s, 3H), 4.03 (m, 4H).

30

Step B: Benzyl 1-[(α -(2-(2-Methyl-1,3-Dioxolan-2-yl)carboxyisobutoxy)carbonyl)aminomethyl]-1-Cyclohexane Acetate (130)

A mixture containing benzyl 1-[(α -chloroisobutoxy)carbonyl]aminomethyl]-1-
cyclohexane acetate (95) (1 g, 2.53 mmol), (129) (673 mg, 5.1 mmol), silver carbonate (557
mg, 2.53 mmol), and triethylamine (709 μ L, 5.1 mmol) in chloroform was stirred at room

temperature for 16 h. After filtration, the filtrate was concentrated. The resulting residue was purified by silica gel chromatography, eluting with 15% ethyl acetate/hexane to afford 510 mg (41%) of the title compound (130). MS (ESI) m/z 492.40 ($M+H^+$).

5 **Step C: 1-[(α -(2-(2-Methyl-1,3-Dioxolan-2-yl)carboxyisobutoxy)carbonyl)-aminomethyl]-1-Cyclohexane Acetic Acid (128)**

A mixture of (130) (470 mg, 0.96 mmol) and 5% Pd-C (catalytic amount) in ethanol was stirred under hydrogen at room temperature for 16 h. Filtration and concentration gave 382 mg (100%) of the title compound (128). 1H NMR ($CDCl_3$, 400 MHz): δ 0.96 (d, J = 6.8 Hz, 3H), 0.97 (d, J = 6.8 Hz, 3H), 1.32-1.58 (m, 10H), 1.59 (s, 3H), 2.06 (m, 1H), 2.32 (s, 2H), 3.26 (m, 2H), 4.08 (m, 4H), 5.29 (t, 1H, NH), 6.55 (d, J = 4.8 Hz, 1H). MS (ESI) m/z 402.32 ($M+H^+$).

The acid form was quantitatively converted to its corresponding sodium salt by dissolution in water (5 mL), addition of an equimolar quantity of 0.5 N $NaHCO_3$, followed by lyophilization.

EXAMPLE 41

In Vitro Determination of Caco-2 Cellular Permeability of Prodrugs

The passive permeability of the prodrugs of the current invention may be assessed *in vitro* using standard methods well known in the art (See, *e.g.*, Stewart, *et al.*, *Pharm. Res.*, 1995, 12, 693). For example, passive permeability may be evaluated by examining the flux of a prodrug across a cultured polarized cell monolayer (*e.g.*, Caco-2 cells). Caco-2 cells obtained from continuous culture (passage less than 28) were seeded at high density onto Transwell polycarbonate filters. Cells were maintained with DMEM/ 10% fetal calf serum + 0.1 mM nonessential amino acids + 2 mM L-Gln, 5% CO_2 / 95% O_2 , 37°C until the day of the experiment. Permeability studies were conducted at pH 6.5 apically (in 50 mM MES buffer containing 1 mM $CaCl_2$, 1mM $MgCl_2$, 150 mM $NaCl$, 3 mM KCl , 1 mM NaH_2PO_4 , 5 mM glucose) and pH 7.4 basolaterally (in Hanks' balanced salt solution containing 10 mM HEPES) in the presence of efflux pump inhibitors (250 μ M MK-571, 250 μ M Verapamil, 1 mM Ofloxacin). Inserts were placed in 12 or 24 well plates containing buffer and incubated for 30 min at 37°C°. Prodrug (200 μ M) was added to the apical or basolateral compartment (donor) and concentrations of prodrug and/or released parent drug in the opposite compartment (receiver) were determined at intervals over 1 hour using LC/MS/MS. Values of apparent permeability (P_{app}) were calculated using the equation:

$$P_{app} = V_r (dC/dt) / (AC_0)$$

Here V_r is the volume of the receiver compartment in mL; dC/dt is the total flux of prodrug and parent drug ($\mu\text{M/s}$), determined from the slope of the plot of concentration in the receiver compartment versus time; C_0 is the initial concentration of prodrug in μM ; A is the surface area of the membrane in cm^2 . Preferably, prodrugs with significant transcellular permeability demonstrate a value of P_{app} of $\geq 1 \times 10^{-6} \text{ cm/s}$ and more preferably, a value of P_{app} of $\geq 1 \times 10^{-5} \text{ cm/s}$, and still more preferably a value of P_{app} of $\geq 5 \times 10^{-5} \text{ cm/s}$. Typical values of P_{app} obtained for prodrugs of GABA analogs are shown in the following table:

Compound	P_{app} (apical to basolateral) (cm/s)	P_{app} (basolateral to apical) (cm/s)	Ratio A-B/B-A
(51)	1.06×10^{-4}	1.25×10^{-5}	8.5
(56)	3.1×10^{-5}	2.0×10^{-6}	15.5
(62)	2.10×10^{-5}	6.40×10^{-6}	3.3
(68)	8.43×10^{-5}	2.26×10^{-5}	3.7
(69)	1.84×10^{-4}	5.22×10^{-6}	35.2
(70)	1.78×10^{-5}	1.68×10^{-6}	10.6
(71)	8.10×10^{-5}	1.99×10^{-5}	4.1
(72)	2.51×10^{-5}	1.26×10^{-6}	2.0
(77)	7.41×10^{-5}	1.43×10^{-5}	5.2
(78)	1.37×10^{-4}	2.46×10^{-5}	5.6
(80)	6.62×10^{-5}	8.75×10^{-6}	7.6
(81)	8.65×10^{-5}	1.27×10^{-5}	6.8
(82)	1.25×10^{-4}	1.82×10^{-5}	6.9
(83)	1.29×10^{-5}	4.48×10^{-5}	0.3
(84)	1.26×10^{-4}	1.57×10^{-5}	8.1
(89)	5.85×10^{-5}	2.34×10^{-6}	25.0
(90)	9.22×10^{-5}	5.75×10^{-6}	16.0

The data in this table shows that the prodrugs disclosed herein have high cellular permeability and should be well absorbed from the intestine. With the exception of compound (83), the apical-to-basolateral permeabilities of these prodrugs exceed their basolateral-to-apical permeabilities. This suggests that these compounds may be substrates for active transport mechanisms present in the apical membrane of Caco cells (though some component of this transcellular permeability may also be mediated by passive diffusion). The greater basolateral-to-apical permeability of (83) suggests that this compound may be subject to efflux across the basolateral membrane, despite the presence of the efflux pump inhibitors MK-571, verapamil and ofloxacin.

EXAMPLE 42

Uptake of Gabapentin Following Administration of Gabapentin or Gabapentin Prodrugs Intracolonicallly in Rats

Sustained release oral dosage forms, which release drug slowly over periods of 6-24 hours, generally release a significant proportion of the dose within the colon. Thus drugs suitable for use in such dosage forms preferably exhibit good colonic absorption. This experiment was conducted to assess the suitability of gabapentin prodrugs for use in an oral sustained release dosage form.

10 **Step A: Administration Protocol**

Rats were obtained commercially and were pre-cannulated in the both the ascending colon and the jugular vein. Animals were conscious at the time of the experiment. All animals were fasted overnight and until 4 hours post-dosing. Gabapentin or gabapentin prodrugs (59), (63), (69), (72), (77), (79), (85), (117) and (126) were administered as a solution (in water or PEG 400) directly into the colon via the cannula at a dose equivalent to 25 mg of gabapentin per kg. Blood samples (0.5 mL) were obtained from the jugular cannula at intervals over 8 hours and were quenched immediately by addition of acetonitrile/methanol to prevent further conversion of the prodrug. Blood samples were analyzed as described below.

Step B: Sample preparation for colonic absorbed drug

1. In blank 1.5 mL eppendorf tubes, 300 μ L of 50/50 acetonitrile/methanol and 20 μ L of *p*-chlorophenylalanine was added as an internal standard.
2. Rat blood was collected at different time points and immediately 100 μ L of blood was added into the eppendorf tube and vortexed to mix.
3. 10 μ L of a gabapentin standard solution (0.04, 0.2, 1, 5, 25, 100 μ g/mL) was added to 90 μ L of blank rat blood to make up a final calibration standard (0.004, 0.02, 0.1, 0.5, 2.5, 10 μ g/mL). Then 300 μ L of 50/50 acetonitrile/methanol was added into each tube followed by 20 μ L of *p*-chlorophenylalanine.
4. Samples were vortexed and centrifuged at 14,000 rpm for 10 min.
5. Supernatant was taken for LC/MS/MS analysis.

Step C: LC/MS/MS analysis

An API 2000 LC/MS/MS spectrometer equipped with Shimadzu 10ADVP binary pumps and a CTC HTS-PAL autosampler were used in the analysis. A Zorbax XDB C8 4.6 x 150 mm column was heated to 45 °C during the analysis. The mobile phase was 0.1% formic acid (A) and acetonitrile with 0.1% formic acid (B). The gradient condition was: 5% B for 1 min, then to 98% B in 3 min, then maintained at 98% B for 2.5 min. The mobile phase was returned to 5%B for 2 min. A TurbolonSpray source was used on the API 2000. The analysis was done in positive ion mode and an MRM transition of 172/137 was used in the analysis of gabapentin (MRM transitions 426/198 for (59), 364/198 for (63), 392/198 for (69), 316/198 for (72), 330/198 for (77), 330/198 for (79), 316/198 for (85) and 327.7/153.8 for (117) were used). 20 µL of the samples were injected. The peaks were integrated using Analyst 1.1 quantitation software. Following colonic administration of each of these prodrugs, the maximum plasma concentrations of gabapentin (C_{max}), as well as the area under the gabapentin plasma concentration vs. time curves (AUC) were significantly greater (> 2-fold) than that produced from colonic administration of gabapentin itself. For example, prodrug (77) provided both gabapentin C_{max} and AUC values greater than 10-fold higher than gabapentin itself. This data demonstrates that compounds of the invention may be formulated as compositions suitable for enhanced absorption and/or effective sustained release of GABA analogs to minimize dosing frequency due to rapid systemic clearance of these GABA analogs.

EXAMPLE 43**Sustained Release of Gabapentin Following Prodrug Administration Using Osmotic Mini-Pump Devices in Beagle Dogs**

Gabapentin or the gabapentin prodrugs (77) and (82) (at a dose equivalent to 10 mg of gabapentin per kg) were dissolved in a suitable solvent (e.g., water, PEG 400, etc.) and filled into preweighed Alzet® mini-osmotic pump devices (Model 2001D) (Durect Corp., Cupertino, CA). The filled Alzets were pre-equilibrated by soaking in isotonic saline at 37°C for 3 hours and stored in sealed containers at 4°C overnight. Alzets were then administered orally to four fasted male beagle dogs (approx. 6.5 kg). Animals were fed at 4 hr after each dose. Blood samples (1.0 mL) were withdrawn at intervals over 48 hours and processed immediately for plasma. Plasma samples were frozen and stored at -80°C until analyzed using the method described above. Both prodrugs afforded gabapentin

concentrations in plasma at 12 hours post-dosing that were greater than 2-fold the concentration of gabapentin seen following administration of gabapentin itself in the Alzet device. This data further confirms that compounds of the invention may be formulated as compositions suitable for effective sustained release of GABA analogs.

5

EXAMPLE 44

Uptake of Pregabalin Following Administration of Pregabalin or Pregabalin Prodrugs **Intracolonicly in Rats**

10 The protocol of Example 41 was repeated with pregabalin and the pregabalin prodrugs (110) and (112). Following colonic administration of each of these prodrugs, the maximum plasma concentrations of pregabalin (C_{max}), as well as the area under the pregabalin plasma concentration vs. time curves (AUC) were significantly greater (> 2-fold) than that produced from colonic administration of pregabalin itself.

15


Finally, it should be noted that there are alternative ways of implementing the present invention. Accordingly, the present embodiments are to be considered as illustrative and not restrictive, and the invention is not to be limited to the details given herein, but may be modified within the scope and equivalents of the appended claims.

20 All publications and patents cited herein are incorporated by reference in their entirety.

CLAIMS


What is claimed is:

1. A compound of Formula (I), Formula (II) or Formula (III):




(1)

5



(II)



(III)

or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:

m, n, t and u are independently 0 or 1;


X is O or NR¹⁶;

W is O or NR¹⁷;

Y is O or S;

R¹ is selected from the group consisting of hydrogen, R²⁴C(O)-, R²⁵OC(O)-,

$$\text{R}^{24}\text{C}(\text{S})-, \text{R}^{25}\text{OC}(\text{S})-, \text{R}^{25}\text{SC}(\text{O})-, \text{R}^{25}\text{SC}(\text{S})-, (\text{R}^9\text{O})(\text{R}^{10}\text{O})\text{P}(\text{O})-, \text{R}^{25}\text{S}-,$$



each R² is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, substituted acyl, acylamino, substituted acylamino, alkylamino, substituted alkylamino, alkylsulfinyl, substituted alkylsulfinyl, alkylsulfonyl, substituted alkylsulfonyl, alkylthio, substituted alkylthio, alkoxycarbonyl, substituted alkoxycarbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, aryloxy, substituted aryloxy, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, dialkylamino, substituted dialkylamino, halo, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl, substituted heteroarylalkyl, heteroalkyloxy, substituted heteroalkyloxy, heteroaryloxy and substituted heteroaryloxy, or optionally, R² and R¹⁶ together with the atoms to which they are attached form a cycloheteroalkyl or substituted cycloheteroalkyl ring;

R³ and R⁶ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl;

R⁴ and R⁵ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, acyl, substituted acyl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroarylalkyl and substituted heteroarylalkyl or optionally, R⁴ and R⁵ together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl or bridged cycloalkyl ring;

R⁸ and R¹² are independently selected from the group consisting of hydrogen, acyl, substituted acyl, alkoxycarbonyl, substituted alkoxycarbonyl, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl, or

optionally R^8 and R^{12} together with the carbon atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring;

R^{11} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, acyl, substituted acyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cyano, cycloalkyl, substituted cycloalkyl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl, substituted heteroarylalkyl, alkoxy carbonyl, substituted alkoxy carbonyl, cycloheteroalkyloxy carbonyl, substituted cycloheteroalkyloxy carbonyl, aryloxy carbonyl, substituted aryloxy carbonyl, heteroaryloxy carbonyl, substituted heteroaryloxy carbonyl and nitro;

R^7 , R^9 , R^{10} , R^{15} , R^{16} and R^{17} are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl;

R^{13} and R^{14} are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy carbonyl, substituted alkoxy carbonyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, carbamoyl, cycloalkyl, substituted cycloalkyl, cycloalkoxy carbonyl, substituted cycloalkoxy carbonyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl or optionally, R^{13} and R^{14} together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring;

R^{20} and R^{21} are independently selected from the group consisting of hydrogen, acyl, substituted acyl, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl or optionally R^{20} and R^{21} together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring;

R^{22} and R^{23} are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl and substituted arylalkyl or optionally, R^{22} and R^{23} together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring;

R^{24} is selected from the group consisting of hydrogen, acyl, substituted acyl, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl,

substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl;

R^{25} is selected from the group consisting of acyl, substituted acyl, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl; and

with the provisos that:

when R^3 and R^6 are both hydrogen, then R^4 and R^5 are neither both hydrogen nor both methyl;

in a compound of Formula (I) when either n is 0 or when n is 1 and X is NR^{16} , then R^1 is not hydrogen;

in a compound of Formula (I) neither R^1 , R^7O- , $R^{24}C(O)-$, $R^{25}C(O)-$ nor $R^{25}O-$ is a moiety derived from a bile acid;

in a compound of Formula (I) when R^1 is $R^{24}C(O)-$ and n is 0, then R^{24} is not methyl, *tert*-butyl, 2-aminoethyl, 3-aminopropyl, benzyl, phenyl or 2-(benzyloxymethyl)phenyl;

in a compound of Formula (I) when R^1 is $R^{25}OC(O)-$, then R^{25} is not $R^{26}C(O)CR^{13}R^{14}-$, wherein R^{26} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl;

in a compound of Formula (I) when R^1 is $R^{25}OC(O)-$ and n is 0, then R^{25} is not methyl, *tert*-butyl or benzyl;


in a compound of Formula (I) when n is 0 and R^1 is $R^{25}C(O)OCR^{13}R^{14}OC(O)-$ then if either R^{13} or R^{14} is hydrogen, alkoxycarbonyl, substituted alkoxycarbonyl, carbamoyl, cycloalkoxycarbonyl or substituted cycloalkoxycarbonyl, the other of R^{13} or R^{14} is not hydrogen;

in a compound of Formula (I) when n is 1, X is NH , R^3 , R^5 and R^6 are each hydrogen, and R^4 is cyclohexyl, then R^2 is not benzyl;

in a compound of Formula (II) when t is 1, u is 0, then neither R^{20} nor R^{21} is 2-hydroxy-3-methyl-5-chlorophenyl; and

in a compound of Formula (II) when u is 1 and X is O, then t is 1.

2. The compound of Claim 1 having Formula (I):




5

3. The compound of Claim 1, wherein R⁷ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl.

4. The compound of Claim 1, wherein Y is O and R⁷ is hydrogen.

5. The compound of Claim 1, wherein Y is O and R⁷ is alkanyl, substituted
15 alkanyl, alkenyl, substituted alkenyl, aryl or substituted aryl.

6. The compound of Claim 5, wherein R⁷ is methyl, ethyl, benzyl, -C(CH₃)=CH₂, -CH₂C(O)N(CH₃)₂,



20 where V is O or CH₂.

7. The compound of Claim 1, wherein n is 0.

8. The compound of Claim 1, wherein n is 1 and X is NH.

9. The compound of Claim 8, wherein R² is selected from the group consisting of hydrogen, alkanyl, substituted alkanyl, aryl, substituted aryl, arylalkanyl, substituted arylalkanyl, cycloalkanyl, heteroarylalkyl and substituted heteroarylalkanyl.

30 10. The compound of Claim 8, wherein R² is selected from the group consisting of hydrogen, alkanyl and cycloalkanyl.

11. The compound of Claim 10, wherein R^2 is selected from the group consisting of hydrogen, methyl, isopropyl, isobutyl, *sec*-butyl, *t*-butyl, cyclopentyl and cyclohexyl.
- 5 12. The compound of Claim 8, wherein R^2 is selected from the group consisting of substituted alkanyl.
13. The compound of Claim 12, wherein R^2 is selected from the group consisting of $-CH_2OH$, $-CH(OH)CH_3$, $-CH_2CO_2H$, $-CH_2CH_2CO_2H$, $-CH_2CONH_2$, $-CH_2CH_2CONH_2$, -
10 $CH_2CH_2SCH_3$, CH_2SH , $-CH_2(CH_2)_3NH_2$ and $-CH_2CH_2CH_2NHC(NH)NH_2$.
14. The compound of Claim 8, wherein R^2 is selected from the group consisting of aryl, arylalkanyl, substituted arylalkanyl and heteroarylalkanyl.
- 15 15. The compound of Claim 14, wherein R^2 is selected from the group consisting of phenyl, benzyl, 4-hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl and 2-indolyl.
16. The compound of Claim 1, wherein n is 1, X is NR^{16} and R^2 and R^{16} together with the atoms to which they are attached form a cycloheteroalkyl or substituted
20 cycloheteroalkyl ring.
17. The compound of Claim 16, wherein R^2 and R^{16} together with the atoms to which they are attached form an azetidine, pyrrolidine or piperidine ring.
- 25 18. The compound of Claim 1, wherein R^3 and R^6 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl and substituted cycloalkyl.
19. The compound of Claim 1, wherein R^3 and R^6 are independently selected
30 from the group consisting of hydrogen and alkanyl.
20. The compound of Claim 1, wherein R^3 and R^6 are both hydrogen.

21. The compound of Claim 1, wherein R^4 and R^5 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroaryl and substituted heteroaryl.

5

22. The compound of Claim 1, wherein R^4 and R^5 are independently selected from the group consisting of hydrogen, alkanyl and substituted alkanyl.

23. The compound of Claim 1, wherein R^4 is hydrogen and R^5 is selected from the group consisting of C_{1-6} alkanyl.

10

24. The compound of Claim 1, wherein R^4 and R^5 together with the carbon atom to which they are attached are cycloalkanyl or substituted cycloalkanyl.

15

25. The compound of Claim 1, wherein R^4 and R^5 together with the carbon atom to which they are attached are selected from the group consisting of cyclobutyl, substituted cyclobutyl, cyclopentyl, substituted cyclopentyl, cyclohexyl and substituted cyclohexyl.


26. The compound of Claim 1, wherein R^4 and R^5 together with the carbon atom to which they are attached are cycloheteroalkyl or substituted cycloheteroalkyl.

20

27. The compound of Claim 1, wherein R^4 and R^5 together with the carbon atom to which they are attached are bridged cycloalkyl.

28. The compound of Claims 1 or 2 derived from a GABA analog of Formula (XIII):

25



(XIII)

wherein the GABA analog of Formula (XIII) is selected from the group consisting of:


30

1-Aminomethyl-1-cyclohexane acetic acid;

1-Aminomethyl-1-(3-methylcyclohexane) acetic acid;

- 1-Aminomethyl-1-(4-methylcyclohexane) acetic acid;
1-Aminomethyl-1-(4-isopropylcyclohexane) acetic acid;
1-Aminomethyl-1-(4-*tert*-butylcyclohexane) acetic acid;
1-Aminomethyl-1-(3,3-dimethylcyclohexane) acetic acid;
5 1-Aminomethyl-1-(3,3,5,5-tetramethylcyclohexane) acetic acid;
1-Aminomethyl-1-cyclopentane acetic acid;
1-Aminomethyl-1-(3-methylcyclopentane) acetic acid;
1-Aminomethyl-1-(3,4-dimethylcyclopentane) acetic acid;
7-Aminomethyl-bicyclo[2.2.1]hept-7-yl acetic acid;
10 9-Aminomethyl-bicyclo[3.3.1]non-9-yl acetic acid;
4-Aminomethyl-4-(tetrahydropyran-4-yl) acetic acid;
3-Aminomethyl-3-(tetrahydropyran-3-yl) acetic acid;
4-Aminomethyl-4-(tetrahydrothiopyran-4-yl) acetic acid;
3-Aminomethyl-3-(tetrahydrothiopyran-3-yl) acetic acid;
15 3-Aminomethyl-5-methyl-hexanoic acid;
3-Aminomethyl-5-methyl-heptanoic acid;
3-Aminomethyl-5-methyl-octanoic acid;
3-Aminomethyl-5-methyl-nonanoic acid;
3-Aminomethyl-5-methyl-decanoic acid;
20 3-Aminomethyl-5-cyclopropyl-hexanoic acid;
3-Aminomethyl-5-cyclobutyl-hexanoic acid;
3-Aminomethyl-5-cyclopentyl-hexanoic acid;
3-Aminomethyl-5-cyclohexyl-hexanoic acid;
3-Aminomethyl-5-phenyl-hexanoic acid;
25 3-Aminomethyl-5-phenyl-pentanoic acid;
3-Aminomethyl-4-cyclobutyl-butyric acid;
3-Aminomethyl-4-cyclopentyl-butyric acid;
3-Aminomethyl-4-cyclohexyl-butyric acid;
3-Aminomethyl-4-phenoxy-butyric acid;
30 3-Aminomethyl-5-phenoxy-hexanoic acid; and
3-Aminomethyl-5-benzylsulfanyl-pentanoic acid.

29. A compound of Formula (IV):



or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:

Y, R², R³, R⁴, R⁵, R⁶, R⁷, R¹³, R¹⁴, R¹⁶, and R²⁵ are as defined in Claim 1;


5 with the provisos that if either R¹³ or R¹⁴ is hydrogen, alkoxycarbonyl, substituted alkoxycarbonyl, carbamoyl, cycloalkoxycarbonyl or substituted cycloalkoxycarbonyl, then the other of R¹³ or R¹⁴ is not hydrogen; and

R²⁵C(O) is not a moiety derived from a bile acid.

10 30. The compound of Claim 29, wherein Y is O and R⁷ is hydrogen.

31. The compound of Claim 29, wherein Y is O and R⁷ is alkanyl, substituted alkanyl, alkenyl, substituted alkenyl, aryl or substituted aryl.

15 32. The compound of Claim 31, wherein R⁷ is methyl, ethyl, benzyl, -C(CH₃)=CH₂, -CH₂C(O)N(CH₃)₂,




where V is O or CH₂.

20 33. The compound of Claim 29, wherein n is 0.

34. The compound of Claim 29, wherein n is 1 and X is NH.

35. A compound of Claim 29 derived from a GABA analog of Formula (XIII):





wherein the GABA analog of Formula (XIII) is selected from the group consisting of:

- 1-Aminomethyl-1-cyclohexane acetic acid;
- 1-Aminomethyl-1-(3-methylcyclohexane) acetic acid;
- 5 1-Aminomethyl-1-(4-methylcyclohexane) acetic acid;
- 1-Aminomethyl-1-(4-isopropylcyclohexane) acetic acid;
- 1-Aminomethyl-1-(4-*tert*-butylcyclohexane) acetic acid;
- 1-Aminomethyl-1-(3,3-dimethylcyclohexane) acetic acid;
- 1-Aminomethyl-1-(3,3,5,5-tetramethylcyclohexane) acetic acid;
- 10 1-Aminomethyl-1-cyclopentane acetic acid;
- 1-Aminomethyl-1-(3-methylcyclopentane) acetic acid;
- 1-Aminomethyl-1-(3,4-dimethylcyclopentane) acetic acid;
- 7-Aminomethyl-bicyclo[2.2.1]hept-7-yl acetic acid;
- 9-Aminomethyl-bicyclo[3.3.1]non-9-yl acetic acid;
- 15 4-Aminomethyl-4-(tetrahydropyran-4-yl) acetic acid;
- 3-Aminomethyl-3-(tetrahydropyran-3-yl) acetic acid;
- 4-Aminomethyl-4-(tetrahydrothiopyran-4-yl) acetic acid;
- 3-Aminomethyl-3-(tetrahydrothiopyran-3-yl) acetic acid;
- 3-Aminomethyl-5-methyl-hexanoic acid;
- 20 3-Aminomethyl-5-methyl-heptanoic acid;
- 3-Aminomethyl-5-methyl-octanoic acid;
- 3-Aminomethyl-5-methyl-nonanoic acid;
- 3-Aminomethyl-5-methyl-decanoic acid;
- 3-Aminomethyl-5-cyclopropyl-hexanoic acid;
- 25 3-Aminomethyl-5-cyclobutyl-hexanoic acid;
- 3-Aminomethyl-5-cyclopentyl-hexanoic acid;
- 3-Aminomethyl-5-cyclohexyl-hexanoic acid;
- 3-Aminomethyl-5-phenyl-hexanoic acid;
- 3-Aminomethyl-5-phenyl-pentanoic acid;
- 30 3-Aminomethyl-4-cyclobutyl-butyric acid;
- 3-Aminomethyl-4-cyclopentyl-butyric acid;
- 3-Aminomethyl-4-cyclohexyl-butyric acid;
- 3-Aminomethyl-4-phenoxy-butyric acid;
- 3-Aminomethyl-5-phenoxy-hexanoic acid; and

3-Aminomethyl-5-benzylsulfanyl-pentanoic acid.

36. A compound of Formulae (VII) or (VIII):



5 or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:


n , R^2 , R^7 , R^{13} , R^{14} , R^{16} and R^{25} are as defined in Claim 1;

with the provisos that if either R^{13} or R^{14} is hydrogen, alkoxycarbonyl, substituted
 10 alkoxycarbonyl, carbamoyl, cycloalkoxycarbonyl or substituted cycloalkoxycarbonyl, then
 the other of R^{13} or R^{14} is not hydrogen; and
 $R^{25}C(O)$ is not a moiety derived from a bile acid.

37. The compound of Claim 36, wherein R^7 is hydrogen.

15 38. The compound of Claim 36, wherein R^7 is alkanyl, substituted alkanyl,
 alkenyl, substituted alkenyl, aryl or substituted aryl.

39. The compound of Claim 38, wherein R^7 is methyl, ethyl, benzyl, -
 $C(CH_3)=CH_2$, $-CH_2C(O)N(CH_3)_2$,



20

where V is O or CH_2 .

40. The compound of Claim 36, wherein n is 0.

41. The compound of Claim 36, wherein n is 1.

42. The compound of Claim 40, wherein R¹³ is methyl, R⁷ and R¹⁴ are hydrogen,
5 and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl,
10 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

15

43. The compound of Claim 40, wherein R¹³ is ethyl, R⁷ and R¹⁴ are hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl,
20 phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

25

44. The compound of Claim 40, wherein R¹³ is propyl, R⁷ and R¹⁴ are hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-

30

yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

5

45. The compound of Claim 40, wherein R^{13} is isopropyl, R^7 and R^{14} are hydrogen, and R^{25} is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

46. The compound of Claim 40, wherein R^{13} is butyl, R^7 and R^{14} are hydrogen, and R^{25} is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

47. The compound of Claim 40, wherein R^{13} is isobutyl, R^7 and R^{14} are hydrogen, and R^{25} is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-

dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

48. The compound of Claim 40, wherein R¹³ is *sec*-butyl, R⁷ and R¹⁴ are hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

49. The compound of Claim 40, wherein R¹³ is *tert*-butyl, R⁷ and R¹⁴ are hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

50. The compound of Claim 40, wherein R¹³ is cyclopentyl, R⁷ and R¹⁴ are hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-

diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

10 51. The compound of Claim 40, wherein R¹³ is cyclohexyl, R⁷ and R¹⁴ are hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

25 52. The compound of Claim 40, wherein R¹³ is phenyl, R⁷ and R¹⁴ are hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

53. The compound of Claim 40, wherein R¹³ is benzyl, R⁷ and R¹⁴ are hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

54. The compound of Claim 40, wherein R¹³ is phenethyl, R⁷ and R¹⁴ are hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

55. The compound of Claim 40, wherein R¹³ is 3-pyridyl, R⁷ and R¹⁴ are hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

56. The compound of Claim 40, wherein R¹³ is methyl, R¹⁴ is methyl, R⁷ is hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

15

57. The compound of Claim 40, wherein R¹³ is methoxycarbonyl, R¹⁴ is methyl, R⁷ is hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

20

25

58. The compound of Claim 40, wherein R¹³ is ethoxycarbonyl, R¹⁴ is methyl, R⁷ is hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl,

30

1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

5

59. The compound of Claim 40, wherein R¹³ is propoxycarbonyl, R¹⁴ is methyl, R⁷ is hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

60. The compound of Claim 40, wherein R¹³ is isopropoxycarbonyl, R¹⁴ is methyl, R⁷ is hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

61. The compound of Claim 40, wherein R¹³ is butoxycarbonyl, R¹⁴ is methyl, R⁷ is hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-

yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

62. The compound of Claim 40, wherein R¹³ is isobutoxycarbonyl, R¹⁴ is methyl, R⁷ is hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

63. The compound of Claim 40, wherein R¹³ is *sec*-butoxycarbonyl, R¹⁴ is methyl, R⁷ is hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

64. The compound of Claim 40, wherein R¹³ is *tert*-butoxycarbonyl, R¹⁴ is methyl, R⁷ is hydrogen, and R²⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-


dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

65. The compound of Claim 40, wherein R^{13} is cyclohexyloxycarbonyl, R^{14} is methyl, R^7 is hydrogen, and R^{25} is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, pentyl, isopentyl, *sec*-pentyl, neopentyl, 1,1-dimethoxyethyl, 1,1-diethoxyethyl, 1-(1,3-dioxolan-2-yl)-ethyl, 1-(1,3-dioxan-2-yl)-ethyl, 1,1-dimethoxypropyl, 1,1-diethoxypropyl, 1-(1,3-dioxolan-2-yl)-propyl, 1-(1,3-dioxan-2-yl)-propyl, 1,1-dimethoxybutyl, 1,1-diethoxybutyl, 1-(1,3-dioxolan-2-yl)-butyl, 1-(1,3-dioxan-2-yl)-butyl, 1,1-dimethoxybenzyl, 1,1-diethoxybenzyl, 1-(1,3-dioxolan-2-yl)-benzyl, 1-(1,3-dioxan-2-yl)-benzyl, 1,1-dimethoxy-2-phenethyl, 1,1-diethoxy-2-phenethyl, 1-(1,3-dioxolan-2-yl)-2-phenethyl, 1-(1,3-dioxan-2-yl)-2-phenethyl, acetyl, propionyl, butyryl, benzoyl, phenacetyl, phenyl, 4-methoxyphenyl, benzyl, phenethyl, styryl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 3-pyridyl.

66. The compound of Claim 41, wherein R^{16} is hydrogen, and R^2 is selected from the group consisting of hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *tert*-butyl, cyclopentyl, cyclohexyl, phenyl, benzyl, 4-hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl, 2-indolyl, $-\text{CH}_2\text{OH}$, $-\text{CH}(\text{OH})\text{CH}_3$, $-\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{SCH}_3$, $-\text{CH}_2\text{SH}$, $-\text{CH}_2(\text{CH}_2)_3\text{NH}_2$ and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NHC}(\text{NH})\text{NH}_2$.

67. The compound of Claim 66, wherein R^7 is hydrogen.

68. The compound of Claim 66, wherein R^7 is methyl, ethyl, benzyl, $-\text{C}(\text{CH}_3)=\text{CH}_2$, $-\text{CH}_2\text{C}(\text{O})\text{N}(\text{CH}_3)_2$,





where V is O or CH₂.

69. The compound of Claim 41, wherein R⁷ is hydrogen, and R² and R¹⁶ together with the atoms to which they are attached form a pyrrolidine ring.


5

70. A compound of Formulae (V) or (VI):



or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:

R¹ is



10

m is 0; and

n, R², R⁷, R⁸, R¹¹, R¹² and R¹⁶ are as defined in Claim 1.

71. The compound of Claim 70, wherein n is 0, R⁷ is hydrogen, R¹ is R¹¹-K-,
 15 where R¹¹ is selected from the group consisting of acetyl, propionyl, butyryl, isobutyryl, cyclohexanecarbonyl, benzoyl, phenacetyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, cyclohexyloxycarbonyl, phenoxycarbonyl, benzyloxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N-propylcarbamoyl, N-isopropylcarbamoyl, N-phenylcarbamoyl, N-benzylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, N,N-dipropylcarbamoyl, N-
 20 pyrrolidinylcarbamoyl, N-piperidinylcarbamoyl and N-morpholinylcarbamoyl; and

K is an alkenylene moiety selected from the group consisting of 1,2-ethenylene, 1,2-propenyl-1-ene, 1,2-butenyl-1-ene, 3-methyl-1,2-butenyl-1-ene, 2-phenyl-1,2-ethenyl-1-ene, 3-phenyl-1,2-propenyl-1-ene, 2-methoxycarbonyl-1,2-ethenyl-1-ene, 2-ethoxycarbonyl-1,2-ethenyl-1-ene, 2-butoxycarbonyl-1,2-ethenyl-1-ene, 1-methyl-1,2-ethenyl-1-ene, 1-methyl-1,2-propenyl-1-ene, 1-methyl-1,2-butenyl-1-ene, 1,3-dimethyl-1,2-butenyl-1-ene, 1-methyl-2-phenyl-1,2-ethenyl-1-ene, 1-methyl-3-phenyl-1,2-propenyl-1-ene, 1-methyl-2-methoxycarbonyl-1,2-ethenyl-1-ene, 1-methyl-2-ethoxycarbonyl-1,2-ethenyl-1-ene, 1-methyl-2-butoxycarbonyl-1,2-ethenyl-1-ene, 1-ethyl-1,2-ethenyl-1-ene, 1-ethyl-1,2-propenyl-1-ene, 1-ethyl-1,2-butenyl-1-ene, 1-ethyl-3-methyl-1,2-butenyl-1-ene, 1-ethyl-2-phenyl-1,2-ethenyl-1-ene, 1-ethyl-3-phenyl-1,2-propenyl-1-ene, 1-ethyl-2-methoxycarbonyl-1,2-ethenyl-1-ene, 1-ethyl-2-ethoxycarbonyl-1,2-ethenyl-1-ene, 1-ethyl-2-butoxycarbonyl-1,2-ethenyl-1-ene, 1-isopropyl-1,2-ethenyl-1-ene, 1-isopropyl-1,2-propenyl-1-ene, 1-isopropyl-1,2-butenyl-1-ene, 1-isopropyl-3-methyl-1,2-butenyl-1-ene, 1-isopropyl-2-phenyl-1,2-ethenyl-1-ene, 1-isopropyl-3-phenyl-1,2-propenyl-1-ene, 1-isopropyl-2-methoxycarbonyl-1,2-ethenyl-1-ene, 1-isopropyl-2-ethoxycarbonyl-1,2-ethenyl-1-ene, 1-isopropyl-2-butoxycarbonyl-1,2-ethenyl-1-ene, 1-phenyl-1,2-ethenyl-1-ene, 1-phenyl-1,2-propenyl-1-ene, 1-phenyl-1,2-butenyl-1-ene, 1-phenyl-3-methyl-1,2-butenyl-1-ene, 1,2-diphenyl-1,2-ethenyl-1-ene, 1,3-diphenyl-1,2-propenyl-1-ene, 1-phenyl-2-methoxycarbonyl-1,2-ethenyl-1-ene, 1-phenyl-2-ethoxycarbonyl-1,2-ethenyl-1-ene, 1-phenyl-2-butoxycarbonyl-1,2-ethenyl-1-ene, 1-benzyl-1,2-ethenyl-1-ene, 1-benzyl-1,2-propenyl-1-ene, 1-benzyl-1,2-butenyl-1-ene, 1-benzyl-3-methyl-1,2-butenyl-1-ene, 1-benzyl-2-phenyl-1,2-ethenyl-1-ene, 1-benzyl-3-phenyl-1,2-propenyl-1-ene, 1-benzyl-2-methoxycarbonyl-1,2-ethenyl-1-ene, 1-benzyl-2-ethoxycarbonyl-1,2-ethenyl-1-ene, 1-benzyl-2-butoxycarbonyl-1,2-ethenyl-1-ene, 1-methoxycarbonyl-1,2-ethenyl-1-ene, 1-methoxycarbonyl-1,2-propenyl-1-ene, 1-methoxycarbonyl-1,2-butenyl-1-ene, 1-methoxycarbonyl-3-methyl-1,2-butenyl-1-ene, 1-methoxycarbonyl-2-phenyl-1,2-ethenyl-1-ene, 1-methoxycarbonyl-3-phenyl-1,2-propenyl-1-ene, 1,2-dimethoxycarbonyl-1,2-ethenyl-1-ene, 1-methoxycarbonyl-2-ethoxycarbonyl-1,2-ethenyl-1-ene, 1-methoxycarbonyl-2-butoxycarbonyl-1,2-ethenyl-1-ene, 1-ethoxycarbonyl-1,2-ethenyl-1-ene, 1-ethoxycarbonyl-1,2-propenyl-1-ene, 1-ethoxycarbonyl-1,2-butenyl-1-ene, 1-ethoxycarbonyl-3-methyl-1,2-butenyl-1-ene, 1-ethoxycarbonyl-2-phenyl-1,2-ethenyl-1-ene, 1-ethoxycarbonyl-3-phenyl-1,2-propenyl-1-ene, 1-ethoxycarbonyl-2-methoxycarbonyl-1,2-ethenyl-1-ene, 1,2-diethoxycarbonyl-1,2-ethenyl-1-ene, 1-ethoxycarbonyl-2-butoxycarbonyl-1,2-ethenyl-1-ene, 1-butoxycarbonyl-1,2-ethenyl-1-ene, 1-butoxycarbonyl-1,2-propenyl-1-ene, 1-

butoxycarbonyl-1,2-butenyl-1-ene, 1-butoxycarbonyl-3-methyl-1,2-butenyl-1-ene, 1-butoxycarbonyl-2-phenyl-1,2-ethenyl-1-ene, 1-butoxycarbonyl-3-phenyl-1,2-propenyl-1-ene, 1-butoxycarbonyl-2-methoxycarbonyl-1,2-ethenyl-1-ene, 1-butoxycarbonyl-2-ethoxycarbonyl-1,2-ethenyl-1-ene and 1,2-dibutoxycarbonyl-1,2-ethenyl-1-ene.

5


72. The compound of Claim 70, wherein n is 1, R¹⁶ is hydrogen, and R² is selected from the group consisting of hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *tert*-butyl, cyclopentyl, cyclohexyl, phenyl, benzyl, 4-hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl, 2-indolyl, -CH₂OH, -CH(OH)CH₃, -CH₂CO₂H, -CH₂CH₂CO₂H, -CH₂CONH₂, -CH₂CH₂CONH₂, -CH₂CH₂SCH₃, -CH₂SH, -CH₂(CH₂)₃NH₂ and -CH₂CH₂CH₂NHC(NH)NH₂.

10

73. The compound of Claim 72, wherein R⁷ is hydrogen.

15

74. The compound of Claim 72, wherein R⁷ is methyl, ethyl, benzyl, -C(CH₃)=CH₂, -CH₂C(O)N(CH₃)₂,



where V is O or CH₂.

20

75. The compound of Claim 70, wherein n is 1, R⁷ is hydrogen, and R² and R¹⁶ together with the atoms to which they are attached form a pyrrolidine ring.

25

76. The compound of Claim 70, wherein n is 0, R⁷ is hydrogen, and R⁸ and R¹² together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring.

30

77. The compound of Claim 76, wherein R⁸ and R¹² together with the carbon atoms to which they are attached form a cyclopent-1-ene, cyclohex-1-ene, 2-cyclopenten-1-one, 2-cyclohexen-1-one, 2-(5H)-furanone or 5,6-dihydro-pyran-2-one ring.

78. The compound of Claim 77, wherein R¹¹ is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl and butoxycarbonyl.

79. The compound of Claim 70, wherein n is 0, R^7 is hydrogen, and R^8 and R^{11} together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl or substituted cycloheteroalkyl ring.



5

80. The compound of Claim 79, wherein R^8 and R^{11} together with the carbon atoms to which they are attached form a γ -butyrolactone, δ -valerolactone or 2,2-dimethyl-1,3-dioxan-4,6-dione ring.


10 81. The compound of Claim 80, wherein R^{12} is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, phenyl, benzyl, methoxycarbonyl, ethoxycarbonyl and butoxycarbonyl.

82. A compound of Formulae (V) or (VI):

15



or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:
 R^1 is



20

and n , R^2 , R^7 , R^{15} and R^{16} are as defined in Claim 1.


83. The compound of Claim 82, wherein n is 0, R⁷ is hydrogen and R¹⁵ selected from the group consisting of methyl, ethyl, propyl, isopropyl, cyclopentyl, cyclohexyl, phenyl, 4-hydroxyphenyl, benzyl, 4-hydroxybenzyl and 3-pyridyl.

5

84. The compound of Claim 82, wherein n is 1, R¹⁶ is hydrogen, and R² is selected from the group consisting of hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *tert*-butyl, cyclopentyl, cyclohexyl, phenyl, benzyl, 4-hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl, 2-indolyl, -CH₂OH, -CH(OH)CH₃, -CH₂CO₂H, -CH₂CH₂CO₂H, -CH₂CONH₂, -
 10 CH₂CH₂CONH₂, -CH₂CH₂SCH₃, -CH₂SH, -CH₂(CH₂)₃NH₂ and -CH₂CH₂CH₂NHC(NH)NH₂.

85. The compound of Claim 84, wherein R⁷ is hydrogen.



15 86. The compound of Claim 84, wherein R⁷ is methyl, ethyl, benzyl, -C(CH₃)=CH₂, -CH₂C(O)N(CH₃)₂,



where V is O or CH₂.

20 87. The compound of Claim 82, wherein n is 1, R⁷ is hydrogen, and R² and R¹⁶ together with the atoms to which they are attached form a pyrrolidine ring.

88. A compound of Claim 1 having the structure of Formulae (IX) or (X):



or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:
 t , R^2 , R^7 , R^{20} and R^{21} are as defined in Claim 1.

5 89. The compound of Claim 88, wherein t is 0.


 90. The compound of Claim 88, wherein t is 1 and R^2 is selected from the group
 consisting of hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *tert*-butyl, cyclopentyl,
 cyclohexyl, phenyl, benzyl, 4-hydroxybenzyl, 4-bromobenzyl, 2-imidazolyl, 2-indolyl, -
 10 CH_2OH , $-\text{CH}(\text{OH})\text{CH}_3$, $-\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CH}_2\text{CONH}_2$, -
 $\text{CH}_2\text{CH}_2\text{SCH}_3$, $-\text{CH}_2\text{SH}$, $-\text{CH}_2(\text{CH}_2)_3\text{NH}_2$ and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NHC}(\text{NH})\text{NH}_2$.

 91. The compound of Claim 89 or 90, wherein R^{20} and R^{21} are independently
 selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl,
 15 heteroaryl and substituted heteroaryl.

 92. The compound of Claim 89 or 90, wherein R^{20} and R^{21} together with the
 carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl,
 cycloheteroalkyl or substituted cycloheteroalkyl ring.


20

 93. The compound of Claims 91 and 92, wherein R^7 is hydrogen, methyl, ethyl,
 benzyl, $-\text{C}(\text{CH}_3)=\text{CH}_2$, $-\text{CH}_2\text{C}(\text{O})\text{N}(\text{CH}_3)_2$,




where V is O or CH₂.

94. A compound of Claim 1 having the structure of Formulae (XI) or (XII):



(XI)



(XII)

- 5 or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:

n, R¹, R², R²² and R²³ are as defined in Claim 1.

95. The compound of Claim 94, wherein n is 1, X is NH, R¹ is hydrogen and R² is arylalkyl.

10

96. The compound of Claim 95, wherein R² is benzyl.

97. The compound of Claim 94, wherein n is 0 and R¹ is R²⁵OC(O)-.

15

98. The compound of Claim 97, wherein R²⁵ is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl.

99. The compound of Claim 98, wherein R²⁵ is ethyl.

20

100. The compound of Claim 94, wherein R²² and R²³ are independently selected from the group consisting of hydrogen, alkyl and substituted alkyl.

101. The compound of any one of Claims 94 – 100, wherein R²² is methyl and R²³ is hydrogen.

25

102. The compound of any one of Claims 94 – 100, wherein R²² is methyl and R²³ is methyl.

103. A method for treating or preventing epilepsy, depression, anxiety, psychosis, faintness attacks, hypokinesia, cranial disorders, neurodegenerative disorders, panic, pain, inflammatory disease, insomnia, gastrointestinal disorders or ethanol withdrawal syndrome in a patient, comprising administering to a patient in need of such treatment a
5 therapeutically effective amount of a compound according to any one of Claims 1, 29, 36, 70, 82, 88 or 94.

104. A method for treating or preventing neuropathic pain, muscular pain or skeletal pain in a patient, comprising administering to a patient in need of such treatment a
10 therapeutically effective amount of a compound according to any one of Claims 1, 29, 36, 70, 82, 88 or 94.

105. A pharmaceutical composition for treating or preventing epilepsy, depression, anxiety, psychosis, faintness attacks, hypokinesia, cranial disorders,
15 neurodegenerative disorders, panic, pain, inflammatory disease, insomnia, gastrointestinal disorders or ethanol withdrawal syndrome in a patient, comprising a therapeutically effective amount of a compound according to any one of Claims 1, 29, 36, 70, 82, 88 or 94 and a pharmaceutically acceptable vehicle.

20 106. A pharmaceutical composition for treating or preventing neuropathic pain, muscular pain or skeletal pain in a patient, comprising a therapeutically effective amount of a compound according to any one of Claims 1, 29, 36, 70, 82, 88 or 94 and a pharmaceutically acceptable vehicle.


25 107. A GABA analog derivative, M-G, for administration to a patient in need of therapy, wherein M is a promoiety and G is derived from a GABA analog, H-G (where H is hydrogen), and where

the promoiety M, once cleaved from G, and any metabolite thereof, exhibits a carcinogenically toxic dose (TD₅₀) in rats of greater than 0.2 mmol/kg/day
30 and the promoiety M is cleaved from G at a sufficient rate *in vivo*, upon colonic administration to rats, to produce:

(i) a maximum concentration of H-G in plasma (C_{max}) of at least 120% of the C_{max} of H-G in plasma achieved by colonically administering an equimolar dose of H-G;
and

(ii) an AUC that is at least 120 % of the AUC achieved by colonically administering an equimolar dose of H-G.

108. A GABA analog derivative, M-G, for administration to a patient in need of therapy, wherein M is a promoiety and G is derived from a GABA analog, H-G (where H is hydrogen), wherein M-G has the structure of Formula (XIV):



(XIV)

or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:

10 Y is O or S;

R is hydrogen, or R and R⁶ together with the atoms to which they are attached form an azetidine, substituted azetidine, pyrrolidine or substituted pyrrolidine ring;

R³ and R⁶ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl;

R⁴ and R⁵ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, acyl, substituted acyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl or optionally, R⁴ and R⁵ together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl or bridged cycloalkyl ring;


R⁷ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, substituted cycloheteroalkyl, heteroalkyl, substituted heteroalkyl, heteroaryl, substituted heteroaryl, heteroarylalkyl and substituted heteroarylalkyl; and

wherein the promoiety M, once cleaved from G, and any metabolite thereof, exhibits a carcinogenically toxic dose (TD₅₀) in rats of greater than 0.2 mmol/kg/day

30 and the promoiety M is cleaved from G at a sufficient rate *in vivo*, upon colonic administration to rats, to produce:

- (i) a maximum concentration of H-G in plasma (C_{\max}) of at least 120% of the C_{\max} of H-G in plasma achieved by colonically administering an equimolar dose of H-G; and
- (ii) an AUC that is at least 120 % of the AUC achieved by colonically administering an equimolar dose of H-G.

109. A GABA analog derivative, M-G, for administration to a patient in need of such therapy, where M is a promoiety and G is derived from a GABA analog, H-G (where H is hydrogen), and wherein M-G has the structure of Formula (XIV), and M has the structure of Formula (XV):




wherein:

n, X, R^1 and R^2 are as previously defined in Claim 1; and

wherein the promoiety M, once cleaved from G, and any metabolite thereof, exhibits a carcinogenically toxic dose (TD_{50}) in rats of greater than 0.2 mmol/kg/day and the promoiety M cleaves from G at a sufficient rate *in vivo*, upon colonic administration to rats, to produce:

- (i) a maximum concentration of H-G in plasma (C_{\max}) of at least 120% of the C_{\max} of H-G in plasma achieved by colonically administering an equimolar dose of H-G; and
- (ii) an AUC that is at least 120 % of the AUC achieved by colonically administering an equimolar dose of H-G.

110. The GABA analog derivative, M-G, of any one of Claims 107 to 109, wherein H-G, once cleaved from M, is substantially free of a lactam having the structure of Formula (XVI):



wherein R is hydrogen.

111. The GABA analog derivative, M-G, of any one of Claims 107-109, wherein
5 M, upon cleavage from G, and any metabolite thereof, is substantially free of formaldehyde.

112. The GABA analog derivative, M-G, of any one of Claims 107-109, wherein
M, upon cleavage from G, and any metabolite thereof, is substantially free of pivalic acid.

10 113. The GABA analog derivative, M-G, of any one of Claims 107-109, wherein
the promoiety M cleaves from G at a sufficient rate *in vivo*, upon colonic administration to
rats, to produce a C_{max} of H-G in plasma of at least 200% of the C_{max} of H-G in plasma
achieved by colonically administering an equimolar dose of H-G.

15 114. The GABA analog derivative, M-G, of any one of Claims 107-109, wherein
the promoiety M cleaves from G at a sufficient rate *in vivo*, upon colonic administration to
rats, to produce a C_{max} of H-G in plasma of at least 1000% of the C_{max} of H-G in plasma
achieved by colonically administering an equimolar dose of H-G.

20 115. The GABA analog derivative, M-G, of any one of Claims 107-109, wherein
the promoiety M cleaves from G at a sufficient rate *in vivo*, upon colonic administration to
rats, to produce an AUC of H-G in plasma of at least 200% of the AUC of H-G in plasma
achieved by colonically administering an equimolar dose of H-G.

25 116. The GABA analog derivative, M-G, of any one of Claims 107-109, wherein
the promoiety M cleaves from G at a sufficient rate *in vivo*, upon colonic administration to
rats, to produce an AUC of H-G in plasma of at least 500% of the AUC of H-G in plasma
achieved by colonically administering an equimolar dose of H-G.

117. The GABA analog derivative, M-G, of any one of Claims 107-109, wherein the promoiety M cleaves from G at a sufficient rate *in vivo*, following oral administration to dogs using a sustained release device at a dose of about 60 μmol equivalent of H-G per kg, to produce a plasma concentration of H-G at 12 hours post dosing of at least 200% of the
- 5 plasma concentration of H-G achieved from an equimolar dose of H-G following the same mode of administration.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 December 2002 (19.12.2002)

PCT

(10) International Publication Number
WO 02/100347 A3

(51) International Patent Classification⁷: **A61K 31/44**,
31/335, 31/27, 31/225, 31/195, A61P 25/00, C07D 213/02,
317/00, C07C 205/00, 229/00, 261/00, 269/00, 271/00

(74) Agents: **SINGH, Sunil, K.**; Cooley Godward LLP 3000
El Camino Real, Five Palo Alto Square, Palo Alto, CA
94306-2155 et al. (US).

(21) International Application Number: PCT/US02/18689

(22) International Filing Date: 11 June 2002 (11.06.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/297,521 11 June 2001 (11.06.2001) US
60/298,514 14 June 2001 (14.06.2001) US
60/366,090 19 March 2002 (19.03.2002) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,
YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

(71) Applicant: **XENOPORT, INC.** [US/US]; 3410 Central
Expressway, Santa Clara, CA 95051 (US).

(72) Inventors: **GALLOP, Mark, A.**; 511 Orange Avenue, Los
Altos, CA 94022 (US). **CUNDY, Kenneth, C.**; 45 Sum-
mit Ridge Place, Redwood City, CA 94062 (US). **ZHOU,**
Cindy, X.; 1054 Colorado Avenue, Palo Alto, CA 94303
(US). **QIU, Fayang, G.**; 5 Hackett Avenue #3, Mountain
View, CA 94043 (US). **YAO, Fenmei**; 2185B Stanford Av-
enue,, Mountain View, CA 94040 (US). **XIANG, Jia-Ning**;
707 DeSoto Drive, Palo Alto, CA 94303 (US). **OLLMAN,**
Ian, R.; 3222 Glendora Drive # 103, San Mateo, CA 94403
(US).

Published:

— with international search report

(88) Date of publication of the international search report:
16 October 2003

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: PRODRUGS OF GABA ANALOGS, COMPOSITIONS AND USES THEREOF

(57) Abstract: The present invention provides prodrugs of GABA analogs, pharmaceutical compositions of prodrugs of GABA analogs and methods for making prodrugs of GABA analogs. The present invention also provides methods for using prodrugs of GABA analogs and methods for using pharmaceutical compositions of prodrugs of GABA analogs for treating or preventing common diseases and/or disorders.

WO 02/100347 A3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/18689

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/44, 31/335, 31/27, 31/225, 31/195; A61P 25/00; C07D 213/02, 317/00; C07C 205/00, 229/00, 261/00, 269/00, 271/00
US CL : 514/356, 467, 478, 547, 561; 546/322; 549/449; 560/115, 125; 562/507

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
U.S. : 514/356, 467, 478, 547, 561; 546/322; 549/449; 560/115, 125; 562/507

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	EP 1 178 034 A1 (WARNER-LAMBERT COMPANY) 06 February 2002, see entire document.	1-116

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "B" earlier application or patent published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

21 February 2003 (21.02.2003)

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Date of mailing of the international search report

14 MAR 2003

Authorized officer
Brenda L. Coleman

Telephone No. 703-308-1235